

Low vitamin D levels may increase risk of Type 1 diabetes

February 4 2013

Having adequate levels of vitamin D during young adulthood may reduce the risk of adult-onset type 1 diabetes by as much as 50%, according to researchers at Harvard School of Public Health (HSPH). The findings, if confirmed in future studies, could lead to a role for vitamin D supplementation in preventing this serious autoimmune disease in adults. The study was published online February 3, 2013 and will appear in the March 1 print edition of the *American Journal of Epidemiology*.

"It is surprising that a serious disease such as [type](#) 1 diabetes could perhaps be prevented by a simple and safe intervention," said lead author Cassandra Munger, research associate in the Department of Nutrition at HSPH.

This study provides the strongest findings to date to suggest that [vitamin D](#) may be protective against type 1 diabetes.

In type 1 diabetes (once called juvenile-onset or insulin-dependent diabetes), the body's [immune system attacks](#) and permanently disables the insulin-making cells in the pancreas. About 5% of the estimated 25.8 million people in the United States with diabetes have type 1, according to the [American Diabetes Association](#). Although it often starts in childhood, about 60% of type 1 diabetes cases occur after age 20.

Previous studies have suggested that a shortage of vitamin D might boost type 1 [diabetes risk](#), although those studies mostly examined the link between vitamin D levels in pregnancy or childhood and the risk of type

1 diabetes in children. Other research, in young adults, uncovered an association between high vitamin D levels and a lowered risk of multiple sclerosis—an autoimmune disease genetically and epidemiologically related to type 1 diabetes—suggesting that inadequate vitamin D in adulthood may be an important risk factor for [autoimmune diseases](#) in general.

Long-term study of military personnel

The researchers conducted a prospective case-control study of U.S. military personnel on active duty, using blood samples from the Department of Defense Serum Repository, which contains more than 40 million samples collected from 8 million military personnel since the mid-1980s. Identifying 310 individuals diagnosed with type 1 diabetes between 1997 and 2009, the team examined blood samples taken before onset of the disease, and compared the samples with those of 613 people in a control group.

The researchers found that white, non-Hispanic, healthy young adults with higher serum levels (>75 nmol/L) of vitamin D had about half the risk of developing type 1 diabetes than those with the lowest levels of vitamin D (

Citation: Low vitamin D levels may increase risk of Type 1 diabetes (2013, February 4) retrieved 26 April 2024 from <https://medicalxpress.com/news/2013-02-vitamin-d-diabetes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.