

Combinations of estrogen-mimicking chemicals found to strongly distort hormone action

March 28 2013

For years, scientists have been concerned about chemicals in the environment that mimic the estrogens found in the body. In study after study, researchers have found links between these "xenoestrogens" and such problems as decreased sperm viability, ovarian dysfunction, neurodevelopmental deficits and obesity. But experimental limitations have prevented them from exploring one of the most serious questions posed by exposure to xenoestrogens: what happens when—as in the real world—an individual is exposed to multiple estrogen-mimicking chemicals at the same time?

Now University of Texas Medical Branch at Galveston researchers have used new techniques to study exposure to low doses of multiple xenoestrogens. And they've come to some disturbing conclusions.

Using <u>cell cultures</u> to test mixtures of three compounds known to affect estrogen signaling,—bisphenol A (found in <u>plastic bottles</u> and the linings), bisphenol S (a supposedly safer replacement for bisphenol A recently found to have similar effects) and nonylphenol (a common component of industrial detergents and surfactants)—the scientists determined that combinations of endocrine disruptors could have a dramatically greater effect than any one of them alone.

"We wanted to see how these persistent, ubiquitous <u>contaminants</u> affect estrogenic signaling when they're mixed together as they are in nature, so



we set up a cell-culture system that allowed us to test their influence on signaling by estradiol, the estrogen found in adult, cycling women," said UTMB professor Cheryl Watson, senior author of a paper on the study now online in the journal *Environmental Health*. "What we found is that these things gang up on estradiol and thwart its response, which is not a good thing."

Watson and her colleagues tested different mixtures of <u>estrogen</u> -disrupting compounds using rat pituitary cells, cells that are master regulators of the animals' endocrine systems. Their experiments measured the responses of key signaling pathways that lead to cell proliferation, the secretion of the pituitary hormone prolactin and the activation of proteins involved in apoptosis (programmed cell death), comparing the effects of estradiol alone with those of estradiol and mixtures of bisphenol A, bisphenol S and nonylphenol.

"These compounds work at very low concentrations—at the parts per trillion or parts per quadrillion level—and when you mix them together they affect estrogenic signaling differently and more dramatically than they do individually," Watson said. "We need to pay attention to this, because estrogens influence so many things in both males and females—reproduction, the immune system, metabolism, bone growth, all sorts of important biological functions."

Studies have detected measurable levels of bisphenol A and bisphenol S in the urine of more than 90 percent of Americans. According to Watson, modern humans are exposed to dozens of <u>xenoestrogens</u> more or less continually.

"These things are all over the environment, and we need to know what they do so we can start figuring out what we need to change," Watson said. "They're probably disrupting and confusing hormones in people, and it's important to find a way to prevent that as soon as we can. We



need to test these compounds for their hormone-disrupting activities before they are put into products, so we can redesign for safety very early in the process."

Provided by University of Texas Medical Branch at Galveston

Citation: Combinations of estrogen-mimicking chemicals found to strongly distort hormone action (2013, March 28) retrieved 3 May 2024 from https://medicalxpress.com/news/2013-03-combinations-estrogen-mimicking-chemicals-stronglydistort.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.