

Stroke risk in elderly treated with antipsychotics is newly linked to specific drug actions

March 11 2013

Antipsychotic administration in the elderly is associated with an increased risk for cerebrovascular accident, more commonly known as stroke; a new study published in *Biological Psychiatry* provides additional insight into this important relationship.

Antipsychotics are prescribed to elderly patients to treat symptoms such as agitation, psychosis, anxiety, insomnia, and depression. The increased risk of stroke associated with these medications was identified approximately a decade ago and has since been replicated by subsequent studies. Although the increase in stroke risk is small, some guidelines discourage the prescription of antipsychotics to elderly patients.

Antipsychotic drugs vary in their effects on the body and so it is likely that antipsychotics are not uniform in their effects on stroke risk. Thus, better understanding of the mechanisms through which antipsychotics increase stroke risk might guide the prescription of safer drugs for elderly patients.

In their study, Wu and colleagues focused on the wide range of brain mechanisms targeted by [antipsychotic medications](#). All antipsychotics block the D2 subtype of receptor for the neurotransmitter dopamine. However, these drugs also act on a range of other receptor targets including the serotonin 5-HT₂ receptor, the M1 receptor for the transmitter acetylcholine, and the alpha₂ receptor for [noradrenaline](#).

To conduct the study, the researchers matched information from a large national database of insurance claims to what is known about the receptor binding profiles of the [antipsychotic drugs](#).

"We found that antipsychotics with a high [binding affinity](#) of alpha2 adrenergic and M1 muscarinic receptors were associated with a greater risk of stroke than the use of other types of antipsychotics," said Dr. Susan Shur-Fen Gau, Professor and Chair at National Taiwan University and corresponding author of the study.

They also found that this stroke risk was elevated in patients who were older and had dementia. Stroke risk was also related to duration and dosage of treatment, with patients who received either short duration of antipsychotic treatment (4 weeks or less) or higher daily doses of treatment showing an increased risk of stroke. This suggests that risk is highest in the initial weeks of antipsychotic treatment and for those with a higher average daily dose.

"Antipsychotics have a wide range of receptor profiles. The [stroke risk](#) profiles from this study suggest that it may be possible to use antipsychotics more safely in the elderly," commented Dr. John Krystal, Editor of [Biological Psychiatry](#).

Additional research is necessary, but the authors agree. They recommend that clinicians start [antipsychotics](#) at low dosages and closely monitoring for side effects in the initial treatment, particularly for individuals with older age and the presence of dementia.

More information: The article is "Association of Stroke with the Receptor-Binding Profiles of Antipsychotics—A Case-Crossover Study" by Chi-Shin Wu, Sheng-Chang Wang, Susan Shur-Fen Gau, Hui-Ju Tsai, and Yu-Cheng Cheng ([doi: 10.1016/j.biopsych.2012.07.006](https://doi.org/10.1016/j.biopsych.2012.07.006)). The article appears in *Biological Psychiatry*, Volume 73, Issue 5 (March 1, 2013)

Provided by Elsevier

Citation: Stroke risk in elderly treated with antipsychotics is newly linked to specific drug actions (2013, March 11) retrieved 9 April 2024 from <https://medicalxpress.com/news/2013-03-elderly-antipsychotics-newly-linked-specific.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.