

Exploring the link between traumatic brain injury and people who are homeless

March 21 2013

Homeless people and their health care providers need to know more about traumatic brain injuries to help prevent and treat such injuries, a new study has found.

[Homeless people](#) have a disproportionately higher risk for TBI compared to the general population, yet little is known about the severity of those injuries, who exactly is suffering from them and what the long-term consequences are.

"A better understanding of TBI, its presentation and characteristics in the [homeless](#) is vital in order to enable appropriate interventions, treatments, and case management in the improvement of outcomes for this important segment of the population," said Dr. Jane Topolovec-Vranic, a clinical researcher in the Trauma and Neurosurgery Program at St. Michael's Hospital.

"Reducing the prevalence of homelessness and the incidence of injury and illness among people who are homeless would have significant financial, societal and individual implications."

Recent research has identified high rates of TBI among homeless people, but there has been no detailed review of existing data. Dr. Topolovec-Vranic and her colleagues in the Head Injury Clinic at St. Michael's Hospital reviewed all the recent scientific studies on homelessness and traumatic brain injuries to identify the gaps in knowledge and suggest areas of future research. Their findings were published recently in the

journal *BMC Public Health*.

As many as one in nine Canadians have experienced homelessness or come close to it. Previous research has found that homeless people often suffer from serious health conditions, use the most expensive [medical interventions](#) such as emergency rooms, require longer hospital stays than people with homes and are at increased risk of death.

The studies Dr. Topolovec-Vranic and her team reviewed found that anywhere between 8 and 53 per cent of homeless people—mostly men—have traumatic brain injuries. The majority suffered a TBI before becoming homeless, suggesting TBI might be a risk factor for homelessness. It's also possible that impulse control disorders, for example, could predispose individuals to both TBI and homelessness, she said.

Traumatic brain injuries are caused by a blow or jolt to the head or a penetrating head injury that disrupts the function of the brain. The severity can range from mild, characterized by a brief change in mental status, to severe, which could include unconsciousness or amnesia. TBI is associated with low subsequent employment rates, which can contribute to a downward spiral into homelessness.

"It is also suggested that in the homeless population, cognitive impairment may increase the risk of remaining homeless, illustrating the potential for TBIs to contribute to the chronicity of homelessness," Dr. Topolovec-Vranic said.

Provided by St. Michael's Hospital

Citation: Exploring the link between traumatic brain injury and people who are homeless (2013, March 21) retrieved 24 April 2024 from <https://medicalxpress.com/news/2013-03-exploring-link->

[traumatic-brain-injury.html](#)

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.