


Intensive glycemic control linked to highest weight gains

March 21 2013



Weight gain is higher in patients with type 2 diabetes mellitus who receive more intensive glycemic control treatment and is associated with a reduction of A1C from baseline, according to research published online Feb. 14 in *Diabetes Care*.

(HealthDay)—Weight gain is higher in patients with type 2 diabetes mellitus (T2DM) who receive more intensive glycemic control treatment and is associated with a reduction of A1C from baseline, according to research published online Feb. 14 in *Diabetes Care*.

Vivian Fonseca, M.D., of Tulane University's Health Sciences Center in New Orleans, and colleagues studied the determinants of weight gain in 8,929 patients with T2DM from the Action to Control Cardiovascular Risk in Diabetes trial who either participated in standard (4,504) or intensive glycemic control treatment arms (4,425).

The researchers found that patients in the intensive glycemia arm gained

an average of 2.7 kg more than those in the standard arm (3.0 versus 0.3 kg). Independent factors associated with weight gain over the two-year treatment duration included younger age, male sex, Asian race, no smoking history, high A1C, baseline [body mass index](#) of 25 to 35 kg/m², high [waist circumference](#), baseline insulin use, and baseline metformin use. Overall, medication use was associated with less than 15 percent of the variability in weight change.

"Following randomization, the intensive group participants with the greatest reduction in A1C gained the most weight," the authors write. "Insulin and TZD [thiazolidinedione] use was associated with the greatest weight gain. Nevertheless, appreciation of these characteristics may help develop strategies to prevent weight gain when initiating intensive glycemic control in the future."

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

[Health News](#) Copyright © 2013 [HealthDay](#). All rights reserved.

Citation: Intensive glycemic control linked to highest weight gains (2013, March 21) retrieved 18 April 2024 from
<https://medicalxpress.com/news/2013-03-intensive-glycemic-linked-highest-weight.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--