

New study finds maternal diet important predictor of severity for infant RSV

March 4 2013

An important predictor of the severity of respiratory syncytial virus (RSV) in infants may be what their mothers ate during pregnancy, according to a Vanderbilt study published in the *American Journal of Respiratory and Critical Care Medicine*.

RSV is the most common cause of severe [lower respiratory tract](#) disease among infants and young children worldwide. Currently there is no effective vaccine against RSV. Outbreaks occur in communities each year, usually lasting 4-5 months during the fall, winter and/or spring months.

Lead author Fernando Polack, M.D., Cesar Milstein Professor of Pediatrics, said his study found that the most serious cases of RSV correlate with mothers who ate a diet high in carbohydrates during pregnancy.

"These cases were not just severe, but the sickest of sick," Polack said. "What we found was the presumptive impact of a carbohydrate-rich diet was clearly dose dependent."

More than 1,200 infants younger than 2 years old were hospitalized in 12 institutions in Buenos Aires, Argentina, during the 2011 RSV season.

Of those patients, nearly 800 were found to have RSV infection, and 106 of those babies were considered to have a life-threatening form of the disease, with [oxygen saturation](#) rates below 87 percent: Twenty-two

infants died in the hospital, and an additional 26 infants died at home with evidence suggesting they died from RSV.

Polack and his collaborators had the mothers fill out a nutrition survey during their final trimester of pregnancy in order to examine the impact of maternal diets heavy in fruits and vegetables versus protein, fats or carbohydrate-rich diets.

Overall, the frequency of life-threatening or fatal RSV in infants was about 12.7 percent among participants, but that rate jumped to 55.6 percent for babies whose mothers ate the greatest proportion of carbohydrates.

"Our study suggests that, where RSV and other [respiratory viruses](#) are concerned, the more sugars a mother eats the worse the situation may get for the baby in the first part of life," said Polack, adding that the next step will be to confirm the association in future studies.

Provided by Vanderbilt University Medical Center

Citation: New study finds maternal diet important predictor of severity for infant RSV (2013, March 4) retrieved 2 May 2024 from <https://medicalxpress.com/news/2013-03-maternal-diet-important-predictor-severity.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--