

# Processed meat linked to premature death

March 6 2013

---

In a huge study of half a million men and women, research in Biomed Central's open access journal *BMC Medicine* demonstrates an association between processed meat and cardiovascular disease and cancer.

One of the difficulties in measuring the effect of eating meat on health is the confounding effect of lifestyle on health. Often vegetarians have healthier lifestyles than the general population, they are less likely to smoke, are less fat, and are more likely to be physically active. Only within a very large study can the consequences of eating meat and [processed meat](#) be isolated from other [lifestyle choices](#).

This EPIC (European Prospective Investigation into Cancer and Nutrition) study involved ten countries and 23 centres in Europe and almost half a million people. In general a diet high in processed meat was linked to other unhealthy choices. Men and women who ate the most processed meat ate the fewest fruit and vegetables and were more likely to smoke. Men who ate a lot of meat also tended to have a high [alcohol consumption](#).

A person's risk of [premature death](#) (increased risk of all cause mortality) increased with the amount of processed meat eaten. This is also true after correcting for confounding variables, although residual confounding cannot be excluded. However, a small amount of red meat appeared to be beneficial which the researchers suggest is because meat is an important source of nutrients and vitamins.

Prof Sabine Rohrmann, from the University of Zurich, who led this

analysis explained, "Risks of dying earlier from cancer and cardiovascular disease also increased with the amount of processed meat eaten. Overall, we estimate that 3% of premature deaths each year could be prevented if people ate less than 20g processed meat per day."

This article marks the launch of an article collection on Medicine for Global Health in *BMC Medicine*. The collection focuses on public health initiatives, the development of health care policies and evidence-based guidelines which are needed to address the global burden of disease. Vulnerable populations, especially in low and middle income countries, continue to be seriously affected by non-communicable and infectious diseases including neglected tropical diseases, while complications during pregnancy and childbirth in these regions leave mothers and infants at risk of severe disability or death.

**More information:** Meat consumption and mortality - results from the European Prospective Investigation into Cancer and Nutrition, Sabine Rohrmann, Kim Overvad, H Bas Bueno-de-Mesquita, Marianne U Jakobsen, Rikke Egeberg, Anne Tjønneland, Laura Nailler, Marie-Christine Boutron-Ruault, Françoise Clavel-Chapelon, Vittorio Krogh, Domenico Palli, Salvatore Panico, Rosario Tumino, Fulvio Ricceri, Manuela M Bergmann, Heiner Boeing, Kuanrong Li, Rudolf Kaaks, Kay-Tee Khaw, Nicholas J Wareham, Francesca L Crowe, Timothy J Key, Androniki Naska, Antonia Trichopoulou, Dimitrios Trichopoulos, Max Leenders, Petra HM Peeters, Dagrun Engeset, Christine Luise Parr, Guri Skeie, Paula Jakszyn, María-José Sánchez, José M Huerta, M Luisa Redondo, Aurelio Barricarte, Pilar Amiano, Isabel Drake, Emily Sonestedt, Göran Hallmans, Ingegerd Johansson, Veronika Fedirko, Isabelle Romieux, Pietro Ferrari, Teresa Norat, Anne C Vergnaud, Elio Riboli and Jakob Linseisen, *BMC Medicine* (in press)

Provided by BioMed Central

Citation: Processed meat linked to premature death (2013, March 6) retrieved 4 May 2024 from <https://medicalxpress.com/news/2013-03-meat-linked-premature-death.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.