

Mortality for acute aortic dissection near one percent per hour during initial onset

March 10 2013

The belief among medical professionals in the 1950s that the mortality rate for type A acute aortic dissection during the initial 24 hours was one to two percent per hour appears to hold true in the contemporary era of treatment, based on a review of the large-scale IRAD registry being presented March 9 at the American College of Cardiology (ACC) Scientific Sessions.

"In the 1950s, the [medical literature](#) suggested that the mortality rate for type A acute aortic dissection was one percent per hour, but we have limited information about the current rate in an era where advanced diagnostic imaging is often readily available," explains lead study author, Craig E. Strauss, MD, MPH, a research cardiologist at the Minneapolis Heart Institute Foundation and physician at the Minneapolis Heart Institute (MHI) at Abbott Northwestern Hospital in Minneapolis. "We reviewed data from the large-scale IRAD registry to obtain a real-world understanding of this patient population in the contemporary era."

The International Registry of Acute Aortic Dissections (IRAD) is a consortium of research centers, including MHI, which evaluates the management and outcomes of acute aortic dissection. Established in 1996, IRAD currently has 30 large referral centers in 11 countries participating in the registry.

"While aortic dissections are a less common diagnosis than a [heart attack](#), [patients](#) present in a similar fashion with significant chest pain or back pain, and the [mortality risk](#) is very real, which is why rapid and

appropriate diagnosis and treatment is so important," says study co-author, Kevin Harris, MD, research [cardiologist](#) at the Minneapolis Heart Institute Foundation.

An aortic dissection is a tear in the wall of the [aorta](#) of the heart that allows blood to flow within the layers of the aorta. Early diagnosis and treatment of this illness is critical for survival. Certain types (type A) of dissections, if left untreated, kill 33% of patients within the first 24 hours, 50% of patients within 48 hours and 75% of patients within two weeks.

For this study, the researchers included 1,969 patients (mean age 61.1, 68.1% male) diagnosed with type A acute aortic dissection in IRAD between January 1996 and February 2012. Of those patients, 195 were managed medically, as opposed to receiving surgery.

The initial 24-hour mortality was higher for the medically managed group, compared with surgically managed patients, based on time from symptom onset (18.5% vs. 3.7%).

The mortality rate per hour during the initial 24 hours was 0.22% for all cases: 0.15% for surgically managed cases and 0.77% for medically managed cases. Among those patients managed medically, the rationale against surgery included age (41.7%), co-morbid illness (64.8%), patient refusal (27.1%), or other (21.3%).

"Essentially, we learned that the mortality rate for medically treated patients with acute [aortic dissection](#) hasn't changed significantly in the past 60 years, despite improvements in pharmacological treatment with beta-blockers and major advancements in [diagnostic imaging](#) tools," says Strauss. CT imaging is the most common method of diagnosis.

"The take-home message from our findings is that mortality is still

extremely high for these patients, which calls for rapid diagnosis and rapid intervention through surgery," adds Harris, who is also co-director of the [Acute Aortic Dissection](#) Program and director of the echocardiography laboratory at the Minneapolis Heart Institute® at Abbott Northwestern Hospital in Minneapolis. "These patients should not be medically managed throughout the night, for instance. If they are an appropriate candidate for surgery, they should be treated immediately."

Provided by Minneapolis Heart Institute Foundation

Citation: Mortality for acute aortic dissection near one percent per hour during initial onset (2013, March 10) retrieved 25 April 2024 from <https://medicalxpress.com/news/2013-03-mortality-acute-aortic-percent-hour.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--