

Mortality higher in Appalachian coal mining counties compared to non-coal mining areas

March 18 2013

West Virginia counties with coal mining activity have higher total mortality rates than their non-coal mining Appalachian counterparts, a University of Pittsburgh Graduate School of Public Health analysis revealed.

"We've known for several years that adverse health outcomes occur at higher rates in Appalachia," said Jeanine Buchanich, Ph.D., deputy director of epidemiology for Pitt Public Health's Center for Occupational Biostatistics and Epidemiology. "For the first time, we've compared Appalachian coal mining counties to non-mining counties matched by median family income and found higher rates of cancer, respiratory disease, diabetes and heart disease in the mining counties."

The findings are reported in the <u>Appalachian Research Initiative for</u> <u>Environmental Science (ARIES) Research Bulletin</u>. Dr. Buchanich will present them in April at the Environmental Considerations in Energy Production symposium in Charleston, W.Va., and they will be concurrently published in the peer-reviewed symposium proceedings published by the Society for Mining, Metallurgy, and Exploration, Inc.

Dr. Buchanich and her colleagues matched 31 West Virginia coal mining counties to non-coal mining counties with comparable family incomes that were still within Appalachia. Some of the non-coal mining matches were outside West Virginia.

The study looked at data on <u>mortality rates</u> related to cancer from 1950



to 2007 and non-cancer deaths from 1960 to 2007.

The data for 2005-2007, the most recent years included, shows that men in coal mining counties had a death rate of 1,200.1 per 100,000, compared to 1,086 per 100,000 in non-coal mining counties. For women the rate was 825.2 per 100,000 for mining counties, compared to 767.2 per 100,000 in non-coal mining counties.

For all types of cancer from 2005-2007, the death rate for men was 266.3 per 100,000 in mining counties, compared to 252 per 100,000 in non-mining counties. For women, it was 180.6 cancer deaths per 100,000 in mining counties and 167.7 per 100,000 in non-mining counties.

"It is interesting that there were higher rates of non-malignant respiratory disease mortality among men, but not women, in coal mining counties," said Dr. Buchanich. "This is indicative of occupational, not environmental, exposures and could reflect lung diseases, such as pneumoconiosis, that are found among people who work in mines."

There were 19 more non-malignant respiratory disease deaths per 100,000 men in mining counties compared to non-mining counties.

Despite the higher overall mortality and cancer death rates in the mining counties, the non-mining counties had higher rates of kidney cancer deaths and stroke.

"The categories where we found excesses in mortality rates in the coal mining counties are arguably heavily influenced by personal behaviors and risk factors, such as smoking, which can cause increased rates of heart disease, diabetes and lung <u>cancer</u>," said Dr. Buchanich. "More studies will be needed to understand the complex interactions of environmental factors, personal behaviors and other risks to determine



the extent coal mining plays in elevating mortality rates."

Provided by University of Pittsburgh

Citation: Mortality higher in Appalachian coal mining counties compared to non-coal mining areas (2013, March 18) retrieved 2 May 2024 from https://medicalxpress.com/news/2013-03-mortality-higher-appalachian-coal-counties.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.