

Neuron loss in schizophrenia and depression could be prevented

March 13 2013

Gamma-aminobutyric acid (GABA) deficits have been implicated in schizophrenia and depression. In schizophrenia, deficits have been particularly well-described for a subtype of GABA neuron, the parvalbumin fast-spiking interneurons. The activity of these neurons is critical for proper cognitive and emotional functioning.

It now appears that parvalbumin [neurons](#) are particularly vulnerable to oxidative stress, a factor that may emerge commonly in development, particularly in the context of [psychiatric disorders](#) like schizophrenia or bipolar disorder, where compromised mitochondrial function plays a role. parvalbumin neurons may be protected from this effect by N-acetylcysteine, also known as Mucomyst, a medication commonly prescribed to protect the liver against the [toxic effects](#) of acetaminophen (Tylenol) overdose, reports a new study in the current issue of *Biological Psychiatry*.

Dr. Kim Do and collaborators, from the Center for Psychiatric Neurosciences of Lausanne University in Switzerland, have worked many years on the hypothesis that one of the causes of schizophrenia is related to vulnerability genes/factors leading to oxidative stress. These oxidative stresses can be due to infections, inflammations, traumas or psychosocial stress occurring during typical [brain development](#), meaning that at-risk subjects are particularly exposed during childhood and adolescence, but not once they reach adulthood.

Their study was performed with mice deficient in glutathione, a

molecule essential for cellular protection against oxidations, leaving their neurons more exposed to the [deleterious effects](#) of oxidative stress.

Under those conditions, they found that the parvalbumin neurons were impaired in the brains of mice that were stressed when they were young. These impairments persisted through their life. Interestingly, the same stresses applied to adults had no effect on their parvalbumin neurons.

Most strikingly, mice treated with the antioxidant N-acetylcysteine, from before birth and onwards, were fully protected against these negative consequences on parvalbumin neurons.

"These data highlight the need to develop novel therapeutic approaches based on antioxidant compounds such as N-acetylcysteine, which could be used preventively in young at-risk subjects," said Do. "To give an antioxidant from childhood on to carriers of a genetic vulnerability for schizophrenia could reduce the risk of emergence of the disease."

"This study raises the possibility that GABA neuronal deficits in psychiatric disorder may be preventable using a drug, N-acetylcysteine, which is quite safe to administer to humans," added Dr. John Krystal, Editor of [Biological Psychiatry](#).

More information: The article is "Early-Life Insults Impair Parvalbumin Interneurons via Oxidative Stress: Reversal by N-Acetylcysteine" by Jan-Harry Cabungcal, Pascal Steullet, Rudolf Kraftsik, Michel Cuenod, and Kim Q. Do ([doi: 10.1016/j.biopsych.2012.09.020](https://doi.org/10.1016/j.biopsych.2012.09.020)). The article appears in *Biological Psychiatry*, Volume 73, Issue 6 (March 15, 2013)

Provided by Elsevier

Citation: Neuron loss in schizophrenia and depression could be prevented (2013, March 13)
retrieved 3 May 2024 from

<https://medicalxpress.com/news/2013-03-neuron-loss-schizophrenia-depression.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.