

Researchers propose a novel prognostic model for disease-specific survival in BCa patients

March 18 2013

A new study from Japan investigated various prognostic indicators, including clinico-pathological and pre-operative hematological factors to develop a novel prognostic factors-based risk stratification model for disease-specific survival (DSS) in bladder cancer (BCa) patients treated with radical cystectomy (RC).

This large series demonstrated that two types of pre-operative haematological disorders assessed by [haemoglobin](#) (Hb) and C-reacted protein (CRP) are independent prognostic indicators for patients with BCa treated by RC.

"Despite the wealth of evidence on the oncological significance of various clinico-pathological and molecular parameters for survival in invasive BCa patients treated with radical cystectomy, alterations of pre-operative haematological parameters have not been sufficiently addressed," write the authors of the study.

"The findings suggest that pre-operative Hb and [CRP levels](#), which can be easily obtained during routine practice, provide additional independent prognostic information for DSS. Moreover, our [risk stratification](#) model, which combines both hematological and pathological parameters, may provide physicians with useful prognostic information for identifying patients who may be candidates for multimodal treatment, including innovative adjuvant treatments."

In the course of the study, the researchers performed a [retrospective analysis](#) of 249 consecutive BCa patients treated with RC without neoadjuvant therapy at several institutions.

Prognostic value of the pre-operative hematological laboratory parameters along with patient's clinico-pathological parameters were evaluated by univariate and multivariate [Cox proportional hazard model](#) analyses. Based on these data, the scientists developed a new scoring model to predict DSS after RC using the regression coefficients of the multivariate model.

One-, 3-, and 5-year DSS rates were 89.1%, 69.3%, and 62.7%, respectively. In multivariate analysis, pre-operative values of low hemoglobin (Hb) (< 0.5 mg / dL), pathological T stage (\geq pT3a), lympho-vascular invasion (LVI) and positive surgical margin (PSM) status were independent factors for predicting poor prognosis. The score was calculated as 2 (if PSM) + 2 (Hb < 0.5 mg / dL) and 0 (if otherwise).

The 4-year DSS in [patients](#) with a score in the ranges 0-1 (low-risk), 2-3 (intermediate-risk) and 4-7 (high-risk) was 83.4%, 48.8% and 5.9%, respectively. All differences in DSS between subgroups were P

More information: T. Sejima, et al., "The prognostic impact of pre-operative hematological disorders and a risk stratification model in bladder cancer patients treated with radical cystectomy," Abstract Nr: 925; 28th Annual EAU Congress, 15 to 19 March 2013; Milan, Italy.

Provided by European Association of Urology

Citation: Researchers propose a novel prognostic model for disease-specific survival in BCa patients (2013, March 18) retrieved 4 May 2024 from

<https://medicalxpress.com/news/2013-03-prognostic-disease-specific-survival-bca-patients.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.