

Resistance to first line anti-malarial drugs is increasing on the Thai-Myanmar border

March 5 2013

Early diagnosis and treatment with antimalarial drugs (ACTs—artemisinin based combination treatments) has been linked to a reduction in malaria in the migrant population living on the Thai-Myanmar border, despite evidence of increasing resistance to ACTs in this location, according to a study by international researchers published in this week's *PLOS Medicine*.

These findings are important as this study suggests that [alternative treatments](#) are urgently needed to replace the failing first line [drug regimen](#) (mefloquine and artesunate).

The authors, led by François Nosten from the Shoklo Malaria Research Unit in Thailand, reached these conclusions by analysing information collected between October 1999 and September 2011 in this region.

They found that the number of confirmed malaria cases with [Plasmodium falciparum](#) (the most serious form of malaria) increased initially, rising from just over 5000 in 2000 to a peak of 13,764 in 2006 and then declining to just over 3,500 in 2011. Encouragingly, the percentage of consultations due to malaria in children aged under 5 years fell from 78% to 7% and the number of new cases of malaria declined from 1.1 to 0.1 episodes per pregnant women-year. In addition, the proportion of patients admitted to hospital with severe disease was stable and the number of deaths from malaria remained extremely low, with an overall case fatality ratio of 0.05%.

The authors also found that the ratio of *P. falciparum* to *P. vivax* (a less severe form of malaria) ratio fell from 1.4 to 0.7 and the rate of *P. falciparum* decreased from 24.3% to 3.4%.

However, worryingly, in the small number of patients undertaking drug resistance tests, the authors found that the effectiveness of ACTs fell steadily with the proportion of patients on treatment but still infected with malaria at day 3, increasing from 0% in 2000 to 28% in 2011.

The authors say: "Despite the emergence of resistance to [ACT] in *P. falciparum*, the strategy of early diagnosis and treatment with ACT has been associated with a reduction in malaria in the [migrant population](#) living on the Thai–Myanmar border."

They continue: "Eliminating *P. falciparum* or dramatically reducing the number of cases is feasible, but the main obstacle is the difficulty of accessing populations living in remote, sometimes dangerous areas and across international borders."

The authors conclude: "Alternative fixed [combination treatments](#) are needed urgently to replace the failing first-line regimen of mefloquine and artesunate."

More information: Carrara VI, Lwin KM, Phyo AP, Ashley E, Wiladphaingern J, et al. (2013) Malaria Burden and Artemisinin Resistance in the Mobile and Migrant Population on the Thai–Myanmar Border, 1999: An Observational Study. *PLoS Med* 10(3): e1001398. [doi:10.1371/journal.pmed.1001398](https://doi.org/10.1371/journal.pmed.1001398)

Provided by Public Library of Science

Citation: Resistance to first line anti-malarial drugs is increasing on the Thai-Myanmar border (2013, March 5) retrieved 26 April 2024 from

<https://medicalxpress.com/news/2013-03-resistance-line-anti-malarial-drugs-thai-myanmar.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.