

Living in a sunny climate does not improve vitamin D levels in hip fracture patients

March 19 2013

While it is well known that a majority of hip fracture patients of all ages and both sexes have insufficient or deficient levels of vitamin D, a new study presented today at the 2013 Annual Meeting of the American Academy of Orthopaedic Surgeons (AAOS) looks at whether or not living in a warm, sunny climate improves patient vitamin D levels.

Researchers retrospectively reviewed the [vitamin D](#) levels of 1,539 patients, including 448 acute hip fracture patients and 1,091 total hip (THR) or total [knee replacement](#) (TKR) patients, from December 2010 to December 2011 at a major medical center in southern California.

Patients were categorized based on their vitamin D levels – deficient, insufficient or sufficient – and according to age and sex. The mean vitamin D levels for the hip fracture and the THR and TKR patients were 26.38 ng/mL and 29.92 ng/mL, respectively. More patients in the hip fracture group were deficient or insufficient (65.8 percent versus 54 percent), and patients age 71 years and older were more deficient or insufficient in the [hip fracture](#) group than the joint replacement group (66.7 percent versus 47.13 percent).

Overall, the majority of patients age 18 and older of both sexes with hip fractures had insufficient levels of vitamin D, and those age 71 or older had significantly lower levels than the [control group](#) of THA and TKA patients.

More information: www.aaos.org/

Provided by American Academy of Orthopaedic Surgeons

Citation: Living in a sunny climate does not improve vitamin D levels in hip fracture patients (2013, March 19) retrieved 5 May 2024 from <https://medicalxpress.com/news/2013-03-sunny-climate-vitamin-d-hip.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.