

Achilles tendon injuries more likely in male 'Weekend Warriors' than others

April 24 2013

Male athletes are the group most likely to tear their Achilles tendon, according to a new study published in the April 2013 issue of *Foot & Ankle International (FAI)*. The activity most likely to cause the injury was basketball, and NBA players such as Kobe Bryant have been in the news lately for this exact injury.

Drs. Steven Raikin, David Garras and Philip Krapchev reviewed 406 records from patients at one clinic diagnosed with [Achilles tendon](#) injuries from August 2000 and December 2010. The average age was 46 years old, 83% of the patients were males, and sports were responsible for 68% of the ruptures.

The most common sports involved were basketball (32% of all ruptures), tennis (9%), and football (8%). Among patients younger than 55 years of age, 77% of ruptures occurred during sports, compared to 42% of the patients 55 or older.

Older [patients](#), and those whose BMI (body-mass index) was greater than 30, were more likely to have non-sports related causes and were more likely to not have been diagnosed correctly at the time of injury. Greater than one-third of the tendon ruptures not caused by sports occurred at work. When the diagnosis was missed, it was usually because the initial diagnosis was an [ankle](#) sprain.

"Delayed diagnosis and treatment have been shown to result in poorer outcomes," says Steven Raikin, MD, of the Rothman Institute in

Philadelphia, PA, and American Orthopaedic [Foot](#) & Ankle Society (AOFAS) member. "Older individuals, and those with a higher BMI, should be evaluated carefully if they have lower leg pain or swelling in the Achilles tendon region."

Re-rupture of the same tendon occurred in 5% of the group, and 6% of the study's population had previously ruptured the other leg's tendon. The study supported previous findings that an Achilles tendon rupture on one leg increases the likelihood of a rupture on the other leg. When the same tendon was re-ruptured, 85% of those injuries had not been treated surgically earlier.

For further information on how to take care of your feet and ankles, or to find a local orthopaedic foot and ankle surgeon, visit the American Orthopaedic Foot & Ankle Society patient website at <http://www.footcaremd.org>. *Foot & Ankle International* is published by SAGE.

More information: "Achilles Tendon Injuries in a United States Population" by Steven M. Raikin, MD; David N. Garras, MD; and Philip V. Krapchev, MD published April 2013 in *Foot & Ankle International*. To read the full text of the article, free for a limited time, click here fai.sagepub.com/content/34/4/475.full

Provided by SAGE Publications

Citation: Achilles tendon injuries more likely in male 'Weekend Warriors' than others (2013, April 24) retrieved 19 April 2024 from <https://medicalxpress.com/news/2013-04-achilles-tendon-injuries-male-weekend.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.