

## 'Mobility shoes' take a load off for knee osteoarthritis sufferers

April 10 2013

---

New research suggests that patients with knee osteoarthritis (OA) who wear flat, flexible footwear (mobility shoes) had significant reduction in knee loading—the force placed upon the joint during daily activities. Results published in *Arthritis & Rheumatism*, a journal of the American College of Rheumatology (ACR), show that long term use of the mobility shoes helped OA patients adapt their gait, or how they walk, which improved knee loading, even when the mobility shoes were no longer worn.

More than 27 million Americans over the age of 25 have some form of OA, which causes painful swelling and stiffness in the hand, foot, [knee](#) or hip joints, according to the ACR. A 2006 study by Hootman et al. published in [Arthritis](#) & Rheumatism projects that doctor-diagnosed arthritis will swell to 67 million U.S. adults by 2030. Furthermore, the Centers for Disease Control and Prevention (CDC) estimate that 16% of adults 45 years of age and older are burdened with symptomatic knee OA.

In their previous studies, Dr. Najia Shakoar and colleagues from Rush University Medical Center in Chicago, Illinois found that walking barefoot as well as with 'mobility [shoes](#),' which are designed to mimic barefoot mechanics, was linked to reduced knee loading compared to when walking with regular footwear worn by participants. However, the authors thought the long term effects of the specialized footwear needed to be evaluated.

Dr. Najia Shakoor explains, "There is much interest in biomechanical interventions, such as orthotic inserts, knee braces, and footwear that aim to improve pain and delay OA progression by decreasing impact on joints. In the present study, we expand understanding of our earlier research by evaluating the impact of the mobility footwear on gait after six months of use." The team recruited 16 participants with knee OA, obtaining a baseline gait with participants walking in their own shoes, mobility shoes, and barefoot. Participants wore the mobility shoes for six hours each day for six days per week and patient gait was evaluated at 6, 12, and 24 weeks in all conditions.

Findings suggest that by 24 weeks participants wearing mobility footwear saw an 18% reduction in knee adduction moment (KAM)—the load on the inner or medial aspect of the knee when walking (where most people develop knee OA) compared to baseline knee loading in their own footwear. No significant difference in KAM was found between walking with mobility shoes and barefoot. Compared to baseline, analyses indicate an 11% and 10% reduction in KAM for OA patients walking in their own shoes and barefoot, respectively, suggesting the mobility shoes may have "re-trained" participant's gait.

"Patients with OA who use flat, flexible footwear may experience a significant reduction in knee loading with continued use," concludes Dr. Shakoor. "Our investigation provides evidence that [footwear](#) choice may be an important consideration in managing knee OA."

**More information:** "Improvement in Knee Loading After Use of Specialized Footwear for Knee Osteoarthritis: Results Of A 6-Month Pilot Investigation." Najia Shakoor, Roy H. Lidtke, Markus A. Wimmer, Rachel A. Mikolaitis, Kharm C. Foucher, Laura E. Thorp, Louis F. Fogg and Joel A. Block. *Arthritis & Rheumatism*; Published Online: April 10, 2013 ([DOI: 10.1002/art.37896](https://doi.org/10.1002/art.37896)).

Provided by Wiley

Citation: 'Mobility shoes' take a load off for knee osteoarthritis sufferers (2013, April 10)  
retrieved 4 April 2024 from  
<https://medicalxpress.com/news/2013-04-mobility-knee-osteoarthritis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.