

# Surgical delay of more than 48 hours increases mortality in older hip fracture patients

April 22 2013

---

Although hip fractures in older patients are known to be a major cause of long term disability and increased risk of death, less is known about the relationship between surgical delay after hip fracture and mortality risk.

A study by Belgian investigators shows that that in older patients with [hip fracture](#), surgical delay of more than 48 hours is significantly and independently associated with increased long-term mortality, even after adjusting for age, sex, and co-morbidities. The study analysed data from 32,383 participants with 1 years of follow-up, extracted from nine clinical cohorts and two population-based databases in the United States, Europe, Israel, Taiwan, and Brazil.

The research was presented (abstract OC18) during the European Congress on Osteoporosis & Osteoarthritis (ESCEO13-IOF) held from April 17 to 20 in Rome, Italy.

Abstracts have been published in '*Osteoporosis International*', Vol. 24, Suppl. 1, 2013

Provided by International Osteoporosis Foundation

Citation: Surgical delay of more than 48 hours increases mortality in older hip fracture patients

(2013, April 22) retrieved 26 April 2024 from <https://medicalxpress.com/news/2013-04-surgical-hours-mortality-older-hip.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.