

Zebrafish study suggests that vitamin B2 (riboflavin) is an antidote to cyanide poisoning

April 30 2013

With the remains of a recent lottery winner having been exhumed for foul play related to cyanide poisoning, future winners might wonder what they can do to avoid the same fate. A new report in [The FASEB Journal](#) involving zebrafish suggests that riboflavin, also known as vitamin B2, may mitigate the toxic effects of this infamous poison. In addition, the report shows that zebrafish are a viable model for investigating the effects of cyanide on humans. As with any research involving animal models, these findings are preliminary until thoroughly tested in clinical trials. Anyone who suspects cyanide poisoning should not attempt to use riboflavin as an antidote, and instead contact local poison control centers or emergency health services immediately.

"We are encouraged to see that many of the effects of cyanide on zebrafish mirror the effects on humans," said Randall Peterson, Ph.D., a researcher involved in the work from the Harvard Medical School in Boston, Massachusetts. "Hopefully, the cyanide biomarkers and antidotes we discover with the help of zebrafish can one day improve our ability to diagnose and treat humans affected by cyanide poisoning."

To make this discovery, scientists exposed zebrafish to cyanide and measured the effects on their behavior, heart rate and survival. The chemical changes that occurred were measured using a [mass spectrometer](#). The effects in zebrafish were then compared to the effects of cyanide on rabbits and humans. Many of the effects in zebrafish

matched those seen in rabbits and humans, confirming that the zebrafish could be used as a model of human cyanide exposure. From there, researchers systematically tested thousands of known drugs to see if any of them could protect the [zebrafish](#) from cyanide toxicity and found four drugs, including riboflavin. The study also identified new biomarkers that indicate cyanide exposure and may be clinically useful for the diagnosis of cyanide exposure and the development of antidotes.

"Lottery winners are not the only ones who have to worry about cyanide poisoning," said Gerald Weissmann, M.D., Editor-in-Chief of *The FASEB Journal*. "Cyanide exposure also occurs through smoke inhalation, industrial accidents, acts of war, and even as an unwanted byproduct of useful drugs. Therefore, the development of antidotes is crucial, and it's nice to know that a likely candidate is already on the drugstore shelf."

More information: Anjali K. Nath, Lee D. Roberts, Yan Liu, Sari B. Mahon, Sonia Kim, Justine H. Ryu, Andreas Werdich, James L. Januzzi, Gerry R. Boss, Gary A. Rockwood, Calum A. MacRae, Matthew Brenner, Robert E. Gerszten, and Randall T. Peterson. Chemical and metabolomic screens identify novel biomarkers and antidotes for cyanide exposure. *FASEB. J.* May 2013 27:1928-1938; [doi:10.1096/fj.12-225037](https://doi.org/10.1096/fj.12-225037)

Provided by Federation of American Societies for Experimental Biology

Citation: Zebrafish study suggests that vitamin B2 (riboflavin) is an antidote to cyanide poisoning (2013, April 30) retrieved 13 March 2024 from <https://medicalxpress.com/news/2013-04-zebrafish-vitamin-b2-riboflavin-antidote.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.