


# Cellular environment controls formation and activity of neuronal connections

May 6 2013

---



Neuron in the net: The illustration shows a neuron from the hippocampus of a mouse in cell culture, which is surrounded by a special structure of the extracellular matrix - a perineuronal net (blue). Various structures of the synapse are coloured red, green and yellow. © RUB, LS Zellmorphologie und

Molekulare Neurobiologie (Department of Cell Morphology and Molecular Neurobiology)

Environment moulds behaviour - and not just that of people in society, but also at the microscopic level. This is because, for their function, neurons are dependent on the cell environment, the so-termed extracellular matrix. Researchers at the Ruhr-Universität have found evidence that this complex network of molecules controls the formation and activity of the neuronal connections. The team led by Dr. Maren Geißler und Prof. Andreas Faissner from the Department of Cell Morphology and Molecular Neurobiology reports in the *Journal of Neuroscience* in collaboration with the team of Dr. Ainhara Aguado, Prof. Christian Wetzel and Prof. Hanns Hatt from the Department of Cell Physiology.

In cooperation with Prof. Uwe Rauch from Lund University in Sweden, Bochum's biologists examined cells from the brains of two mouse species: a species with a normal extracellular matrix and a species which lacked four components of the extracellular matrix due to [genetic manipulation](#), namely the molecules tenascin-C, tenascin-R, neurocan and brevican. They took the cells from the hippocampus, a [brain structure](#) that is crucial for the long-term memory. The team not only examined neurons but also astrocytes, which are in close contact with the neurons, support their function and secrete molecules for the extracellular matrix.

## **Formation, stability and activity of the neuronal connections depend on the matrix**

The researchers cultivated the neurons and astrocytes together for four weeks with a specially developed culture strategy. Among other things,

they observed how many connections, known as synapses, the neurons formed with each other and how stable these were over time. If either the [astrocytes](#) or the neurons in the culture dish derived from animals with a reduced extracellular matrix, these synapses proved to be less stable in the medium term, and their number was significantly reduced. Together with the Department of Cell Physiology at the RUB and the University of Regensburg, the team also showed that the neurons with a mutated matrix showed lower spontaneous activity than normal cells. The extracellular matrix thus regulates the formation, stability and activity of the [neuronal connections](#). The researchers also examined a special structure of the extracellular matrix, the so-called perineuronal nets, which the Nobel laureate Camillo Golgi first described more than a century ago. They were significantly reduced in the environment of genetically modified cells.

**More information:** M. Geissler, C. Gottschling, A. Aguado, U. Rauch, C.H. Wetzel, H. Hatt, A. Faissner (2013): Primary hippocampal neurons, which lack four crucial extracellular matrix molecules, display abnormalities of synaptic structure and function and severe deficits in perineuronal net formation, *Journal of Neuroscience*, [DOI: 10.1523/JNEUROSCI.3275-12.2013](#)

Provided by Ruhr-Universitaet-Bochum

Citation: Cellular environment controls formation and activity of neuronal connections (2013, May 6) retrieved 23 April 2024 from <https://medicalxpress.com/news/2013-05-cellular-environment-formation-neuronal.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.