

Consumers largely underestimating calorie content of fast food

May 23 2013

People eating at fast food restaurants largely underestimate the calorie content of meals, especially large ones, according to a paper published today in *BMJ*.

From 2006 to 2010 many American states and cities passed laws requiring chain restaurants to print [calorie content](#) on menus. The US [Patient Protection](#) and Affordable Care Act of 2010 included a provision that will require all restaurant chains with more than 20 US sites to print calorie content on menus.

Previous research has shown that adults and children underestimate calorie content often by large amounts. However, these studies did not monitor consumer choices at restaurants or focused on a narrow range of [fast food restaurants](#) or individuals.

Researchers therefore carried out a large cross sectional study of repeated visits to fast food restaurant chains in 89 restaurants across the New England region of America: McDonald's, [Burger King](#), Wendy's, KFC, Subway and Dunkin' Donuts. At the time of data collection, none of the chains routinely printed calorie content on menus.

Researchers enrolled a diverse group of adults (>18 years old), [adolescents](#) (aged 11-20) and [school age](#) children (3-15 years old). They collected receipts from participants to calculate the calorie content of their meals, and they administered a short questionnaire which included a question asking participants to estimate the calorie content of their

meal. Parents provided answers for the school age children. The final sample size was 1877 adults, 1178 adolescents and 330 school age children.

Among adult participants, 65% were either overweight or obese, as were 34% of adolescents and 57% of school age children. Less than one quarter of participants reported noticing calorie information in the restaurants, and less than 5% reported using it to help them choose their meal.

The mean calorie content of meals was 836 for adults, 756 for adolescents and 733 for children. On average, adults, adolescents, and parents of school-age children underestimated calorie content by 175 calories, 259 calories and 175 calories, respectively. Two thirds of all participants underestimated the calorie content of their meals with approximately one quarter underestimating the calorie content by at least 500 calories. Participants consuming high calorie meals underestimated by a greater amount than small calorie meals. Underestimation was greater among Subway diners than at any other chain. Adults and adolescents eating at Subway underestimated by 20% and 25% more than those at McDonald's.

The researchers conclude that adults, adolescents and parents of school age children generally underestimate the calorie content of meals, especially large meals and when eating at Subway restaurants. They say that the "forthcoming calorie menu labelling requirements of the US Patient Protection and [Affordable Care](#) Act might help to correct underestimation of calorie content".

The researchers suggest that in addition to providing calories on menus, policymakers could perhaps improve menu labelling by supporting social marketing campaigns to better explain the concept of calories.

More information: Consumers' estimation of calorie content at fast food restaurants: cross sectional observational study, *BMJ*, 2013.

Provided by British Medical Journal

Citation: Consumers largely underestimating calorie content of fast food (2013, May 23)
retrieved 20 March 2024 from <https://medicalxpress.com/news/2013-05-consumers-largely-underestimating-calorie-content.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--