

## **Discovery reshapes understanding of embryonic development**

May 1 2013, by Juliana Bunim



(Medical Xpress)—When a baby is born, one of the first things a parent will do is count fingers and toes. New research conducted at UC San Francisco sheds lights on how these specialized digits are formed.

In a finding likely to fundamentally reshape biologists' understanding of how cells communicate to each other during development, researchers have found a new type of <u>cellular structure</u> that passes along instructions critical to the formation of the limbs and digits of developing embryos.



As a national leader in <u>neonatology</u>, UCSF's mission is to initiate and rigorously pursue both basic science and clinical research to better treat, and one day eliminate, neonatal diseases. UCSF Benioff Children's Hospital is one of the top ranked children's hospitals in the country, and is committed to training physician-scientists to investigate human development and the root causes of congenital anomalies.

"The findings are a significant advance in developmental biology that will help to understand the basis of certain birth defects that alter the number or specification of fingers and toes," said David Rowitch, MD, PhD, chief of neonatology at the children's hospital.

The research was published online April 28 in Nature.

The researchers focused on the molecule called "Sonic hedgehog" (SHH), which acts to pattern <u>precursor cells</u> that make the specialized digits in a developing chick embryo. Equipped with custom-made, highpowered microscopes to visualize the cells and an approach that avoided damaging tissue preparation, the researchers got an unprecedented look at previously invisible extensions of cells that form tube-like projections.

"The presence of these structures, which we call filopodia, was really unexpected," said senior author Maria Barna, PhD, who conducted the research at UCSF and is now an assistant professor of <u>developmental</u> <u>biology</u> and of genetics at Stanford University.

## The role of filopodia

Filopodia have never been reported in vertebrates before, although they were suggested by previous studies in the fruit fly drosophila by UCSF investigator Tom Kornberg, MD, of the UCSF Cardiovascular Research Institute.



The researchers found that the newly identified, nearly invisible fingerlike projections burrow through tightly packed tissue in developing chick and mouse embryos for several cell-lengths. Once contact between cells through these structures has been made, signaling molecules can be delivered directly from one cell to another.

The researchers believe that <u>cells</u> use specialized filopodia as a means of targeted delivery of key signaling molecules in the limb to instruct formation of, say, a thumb versus a pinky finger. If that communication mechanism breaks down, then malformations occur – such as too many or too few fingers or toes.

The wider implication for this research is that similar mechanisms might underlie developmental pattern formation in other organs such as the brain. If so, scientists can learn how and why congenital anomalies occur in other parts of the body.

More information: <u>www.nature.com/nature/journal/ ...</u> <u>ull/nature12157.html</u>

Provided by University of California, San Francisco

Citation: Discovery reshapes understanding of embryonic development (2013, May 1) retrieved 22 May 2024 from <u>https://medicalxpress.com/news/2013-05-discovery-reshapes-embryonic.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.