

Heart failure accelerates male 'menopause'

May 25 2013

Heart failure accelerates the aging process and brings on early andropausal syndrome (AS), according to research presented today at the Heart Failure Congress 2013. AS, also referred to as male 'menopause', was four times more common in men with heart failure.

The <u>Heart Failure</u> Congress is the main annual meeting of the Heart Failure Association of the <u>European Society of Cardiology</u> and is being held 25-28 May in Lisbon, Portugal.

As <u>men</u> get older they are more likely to suffer from andropausal syndrome (AS), also called 'menopause', androgen deficiency in the aging male (ADAM), or late-onset hypogonadism. Men with AS have decreased levels of anabolic hormones, including testosterone, and it has been suggested that these hormone deficiencies are what cause the clinical symptoms.

The symptoms of AS according to the Aging Male Symptom Rating Scale can be divided into three categories: sexual (erectile dysfunction, problems with <u>libido</u>, decrease in beard growth, feelings of 'having passed the zenith of life'), psychological (feeling discouraged, depressed, irritable, anxious, nervous), and somato-vegetative (joint and muscle complaints, sweating, need for more sleep, <u>sleep disturbances</u>, weakness, exhaustion).

Heart failure increases with age. Deficiencies of anabolic hormones are common in men with systolic heart failure, leading to reduced <u>exercise</u> <u>capacity</u>, depression and <u>poor prognosis</u>. But until now the impact of



heart failure on the prevalence of AS and the severity of andropausal symptoms has not been studied.

Professor Ewa A. Jankowska (Wroclaw, Poland) said: "AS leads to poor quality of life. We wanted to discover whether heart failure increases AS and whether additional androgen therapies could improve quality of life in <u>heart failure patients</u>."

For the study, the researchers compared the prevalence of AS and the severity of andropausal symptoms between 232 men with systolic heart failure aged 40-80 years and 362 age-matched healthy peers. The magnitude of andropausal symptoms (psychological, sexual and somato-vegetative) was assessed using the Aging Males' Symptoms (AMS) Rating Scale and AS was diagnosed if the total AMS score was 50 points or more.

They found that AS affected almost one-third of men with heart failure, regardless of their age group. In men aged 40-59 years, heart failure led to a four-fold increase in the prevalence of AS (28% vs. 7%, p

Citation: Heart failure accelerates male 'menopause' (2013, May 25) retrieved 5 May 2024 from <u>https://medicalxpress.com/news/2013-05-heart-failure-male-menopause.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.