

Local community group activities may help reduce neonatal mortality in Vietnam

May 14 2013

Community groups in rural Vietnam comprised of local health workers, politicians and laywomen (Maternal and Newborn Health Groups) set up to tackle challenges to maternal and neonatal health may reduce the neonatal death rate after three years and increase antenatal care attendance, according to a study by researchers from Sweden and Vietnam published in this week's *PLOS Medicine*.

These findings are important as they show that is it feasible to implement community-based activity into the public sector system at low cost.

The researchers (also the authors of this paper) led by Lars-Åke Persson from Uppsala University in Sweden reached these conclusions by conducting a cluster randomized trial involving 90 local communities (communes) in the Quang Ninh province of northeast of Vietnam between 2008 and 2011. In 44 communes, Maternal and [Newborn Health](#) Groups were established (the intervention) by recruiting local [health workers](#), politicians, and women's union representatives to identify and prioritize local perinatal health problems and implement actions to help overcome these problems.

The authors found that the number of babies dying (neonatal mortality rate) did not reduce over the first two years of the intervention but there was a fall in [death rates](#) in the third year to half that observed in the control communes—odds ratio 0.51. Women in the intervention communes were also twice as likely to attend antenatal care than women

in the control group—odds ratio 2.27.

However, the authors also found that there were no differences between the intervention and control communes in other important outcomes, including tetanus immunization, delivery preparedness, institutional delivery, temperature control at delivery, early initiation of breastfeeding, or home visit of a midwife during the first week after delivery.

The authors conclude: "A randomized facilitation intervention with local stakeholder groups composed of primary care staff and local politicians working for three years with a perinatal problem-solving approach resulted in increased attendance to antenatal care and reduced neonatal mortality after a latent period."

They add: "We have shown that community-based participatory approaches to reduce [neonatal mortality](#) is not only effective in South Asian societies but also in a South-East Asian society in rapid transition."

More information: Persson LÅ, Nga NT, Målqvist M, Thi Phuong Hoa D, Eriksson L, et al. (2013) Effect of Facilitation of Local Maternal-and-Newborn Stakeholder Groups on Neonatal Mortality: Cluster-Randomized Trial. PLoS Med 10(5): e1001445.
[doi:10.1371/journal.pmed.1001445](https://doi.org/10.1371/journal.pmed.1001445)

Provided by Uppsala University

Citation: Local community group activities may help reduce neonatal mortality in Vietnam (2013, May 14) retrieved 4 May 2024 from <https://medicalxpress.com/news/2013-05-local-group-neonatal-mortality-vietnam.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.