

Nerve stimulation helps with overactive bladder

May 30 2013



Electric impulses from a hand-held device travel from the nerves in the ankle to the nerves that control pelvic floor function.

Beaumont Health System research finds that symptoms of overactive bladder, or OAB, were reduced in those who received tibial nerve stimulation. The three-year results published in the June issue of *The Journal of Urology* show participants with urinary frequency, urgency and involuntary loss of urine maintained significant improvement in their symptoms.

Tibial [nerve stimulation](#) is a painless procedure that takes place in an outpatient setting. A slim needle [electrode](#) is inserted in the ankle, near the tibial nerve. It carries electric impulses from a hand-held stimulator

to the nerves in the spinal cord that control [pelvic floor](#) function.

Principal investigator Kenneth Peters, M.D., chief of Urology at Beaumont Hospital, Royal Oak and a team of researchers reviewed data of 29 patients who initially responded to 12 weekly neuromodulation system treatments for OAB. [Study participants](#) were followed for three years.

Participants received an average of one tibial nerve treatment per month. After 14 weeks of treatment, 77 percent of patients maintained "moderate or marked improvement" in OAB symptoms.

For those who participated in the study, results show frequent trips to the bathroom during the day decreased by nearly 30 percent, or from 12 to 8.7; nighttime trips decreased by almost 40 percent, or from 2.7 to 1.7; and urge incontinence episodes per day decreased by 100 percent.

"This study demonstrates that with ongoing therapy patients with [overactive bladder](#) can have fewer symptoms and can return to daily activity without disruption or embarrassment that is often caused by this condition," says Dr. Peters.

According to the Urology Care Foundation, about 33 million Americans – men and women – have OAB. The number of people diagnosed with OAB may be much larger because many people living with this condition don't ask for help; they are embarrassed or unaware of available treatments.

More information: [www.jurology.com/article/S0022 ... \(12\)05807-7/abstract](http://www.jurology.com/article/S0022-(12)05807-7/abstract)

Provided by Beaumont Health System

Citation: Nerve stimulation helps with overactive bladder (2013, May 30) retrieved 6 May 2024 from <https://medicalxpress.com/news/2013-05-nerve-overactive-bladder.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.