

Rethinking treatment goals improves results for those with persistent anorexia

May 8 2013

A new, multinational randomized clinical trial has found that patients with severe and enduring anorexia nervosa will not only stick with treatments but also make significant improvements with just a slight modification of the standard goals and methods of treatment.

More than 85 percent of those who enrolled in the trial completed treatment—almost three times the usual retention rate. After eight months of outpatient treatment, patients in both treatment groups reported improved quality of life, reduced symptoms of [mood disorders](#) and enhanced social adjustment.

One crucial element of this trial was the decision to reprioritize how the goals of treatment were presented to those enrolled in this study. Treatment of newly diagnosed adolescents and [young adults](#) tends to focus on weight recovery. For this study, conducted at the University of Sydney, University of London and University of Chicago, treatment goals were set collaboratively, by the care team and the patient, with less emphasis on [weight gain](#) and more on quality of life, reduction of mood disorders, and enhanced social adjustment.

"Our goal was to peel back the [negative impact](#) of anorexia and to shift the traditional pressure to gain weight to an emphasis on improving quality of life and overall functioning," said the study's lead author Stephen Touyz, PhD, from the Centre for Eating and Dieting Disorders at the University of Sydney. "By refocusing the core of treatment, we were able to engage highly resistant individuals with severe and enduring

[anorexia nervosa](#) in treatment, circumvent the notoriously high dropout rates and help them make their lives a little better."

Patients with severe and enduring anorexia nervosa (SE-AN) are notoriously difficult to treat. Most of them develop anorexia as [adolescents](#) and do not respond to treatment. Participants in this study had suffered from severe anorexia for at least seven years, with an average duration of illness of more than 15 years.

Anorexia nervosa has the highest mortality rate of any psychiatric disorder. Most individuals with SE-AN have been through multiple unsuccessful treatment episodes. Repeated failure leaves them with low motivation and a self image dominated by illness.

After prolonged malnourishment, many of those with SE-AN are disabled and unable to hold a job. They often suffer from neurocognitive, cardiac or liver problems, as well as osteoporosis. Insurance companies frequently refuse to pay for treatment because so few individuals improve.

This study, published online by the journal *Psychological Medicine* on May 3, 2013, was the first [randomized clinical trial](#) for chronic anorexia nervosa. It compared two standard treatments but modified them in ways that made them more suitable for individuals with such a profound and persistent disorder. The treatments—cognitive behavioral therapy (CBT) and specialist supportive clinical management (SSCM)—were altered to focus on retention, improved quality of life and to avoid "further failure experiences."

The researchers enrolled 63 patients, all female, at two clinical centers: at the University of Sydney and St George's Hospital, University of London. The average body mass index of study participants was 16.2, with a range from 11.8 to 18.5. (Normal BMI ranges from 18.5 to 25.)

The University of Chicago served as the data-coordinating center.

Thirty-one patients were assigned to CBT and 32 to SSCM. Both treatments involved 30 outpatient therapy sessions over eight months. Patients were assessed at the end of treatment, with follow-up assessments six and 12 months later.

Results from both treatments were comparable, with significant improvements ranging from "moderate" to "large" on many measures. CBT had a greater impact on eating-disorder symptoms and readiness to change. SSCM produced larger improvements in health-related quality of life and depression. Average BMI for both groups increased from 16.2 to 16.8, about 4 percent.

"The results were far better than most people in the field would have expected," said Daniel Le Grange, PhD, professor of psychiatry and director of the eating disorders program at the University of Chicago and the principal investigator for the data-coordinating center. "Many of these patients were profoundly ill. The prevailing wisdom is that current treatments have not been effective and patients are best served by refeeding in the hospital setting. This study showed that specific modification of these behavioral approaches could overcome the high dropout rates and lead to meaningful positive change."

Limitations of the study include a moderate sample size, 63 study participants, and a relatively short follow-up period, 12 months for a disorder that had been present for more than seven years. Despite the limitations, the high retention rate and the magnitude of improvement in most of the outcome measures were very encouraging, especially in the setting of a disorder with poor compliance and limited previous clinical success.

"This study clearly shows that SE-AN patients do respond to, and benefit

from, two specialized treatments when done by clinicians with specialist knowledge," the authors wrote. "This study should provide hope for those suffering from severe and enduring AN as well as stimulate interest in the development of new psychosocial [treatment](#) approaches."

More information: The paper, "Treating severe and enduring anorexia nervosa: a randomized clinical trial," can be found at:
[dx.doi.org/10.1017/S0033291713000949](https://doi.org/10.1017/S0033291713000949).

Provided by University of Chicago Medical Center

Citation: Rethinking treatment goals improves results for those with persistent anorexia (2013, May 8) retrieved 13 March 2024 from <https://medicalxpress.com/news/2013-05-rethinking-treatment-goals-results-persistent.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--