

Weight of tissue flaps affects complication risk after breast reconstruction

May 29 2013

For women undergoing breast reconstruction after mastectomy, the weight of the tissue flap used affects the risk of an important complication called fat necrosis, reports a study in [Plastic and Reconstruction Surgery-Global Open](#).

Information on flap weight—and the number of blood vessels supplying the flap—can help plastic and reconstructive surgeons optimize outcomes for women undergoing [breast reconstruction](#) after mastectomy for breast cancer, according to the study by ASPS Member Surgeon Dr. Justin M. Sacks and colleagues from the Department of [Plastic and Reconstructive Surgery](#) at the Johns Hopkins School of Medicine, Baltimore.

Flap Weight and Blood Supply Predict Fat Necrosis Risk

The researchers analyzed [risk factors](#) for fat necrosis in 123 women undergoing [breast reconstruction](#) after [mastectomy](#) for [breast cancer](#). Fat necrosis refers to the death of [fat cells](#) in the reconstructed breast; it is caused by problems with blood supply to the transplanted flap.

The reconstructions were performed using the women's own tissues—specifically, using flaps of tissue transferred from the abdominal area. In performing autologous reconstructions, plastic and reconstructive surgeons create custom-designed flaps of muscle, fat and

skin from the abdominal area, along with their supplying blood vessels.

Dr. Sacks and coauthors evaluated surgical and patient-related factors affecting the risk of fat necrosis after reconstruction—including the total weight of the abdominal tissue flap before it was transferred to the chest. The analysis included a total of 179 flaps (including 56 women undergoing reconstruction of both breasts).

Fat necrosis occurred in 7.5 percent of flaps overall. The risk was significantly higher for women with heavier tissue flaps: for each 100-gram increase in flap weight, the risk of fat necrosis increased by 50 percent. (The average flap weight was approximately 650 grams.)

Risk of fat necrosis was also affected by the number of supplying blood vessels, or "perforators"—especially in heavier flaps. Among flaps larger than 1,000 grams, the rate of fat necrosis was about 43 percent in those with a single perforator, compared to 14 percent for those with two or more perforators.

Attention to Risk Factors May Improve Breast Reconstruction Outcomes

African American women were also at higher risk of fat necrosis after breast reconstruction, with a risk nearly 12 times higher than in Caucasian women. Risk was not significantly affected by patient age, body weight, and various medical and surgical factors.

Fat necrosis is an important complication after breast reconstruction. "Development of fat necrosis requiring reoperation can result in significant aesthetic deformity as well as emotional and financial costs," Dr. Sacks and coauthors note.

Plastic and reconstructive surgeons are constantly working to refine the technical aspects of breast reconstruction, including flap design, in order to provide the best possible surgical and aesthetic results. Although other studies have looked at risk factors for fat necrosis, the new study is the first to focus on the flap weight and its interaction with blood supply.

The results show that larger, heavier flaps are at higher risk of fat necrosis, particularly in flaps with just one perforating blood vessel. Dr. Sacks and colleague suggest that plastic and reconstructive surgeons should consider "balancing perforator number and increasing flap weight" in designing abdominal flaps for autologous breast reconstruction. They suggest to ensure adequate [blood supply](#) to flaps in African American women, who appear to be at higher risk of fat necrosis after breast reconstruction.

Provided by Wolters Kluwer Health

Citation: Weight of tissue flaps affects complication risk after breast reconstruction (2013, May 29) retrieved 20 April 2024 from <https://medicalxpress.com/news/2013-05-weight-tissue-affects-complication-breast.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--