

Alzheimer's and low blood sugar in diabetes may trigger a vicious cycle

June 10 2013

Diabetes-associated episodes of low blood sugar may increase the risk of developing dementia, while having dementia or even milder forms of cognitive impairment may increase the risk of experiencing low blood sugar, according to a UC San Francisco scientist who led a new study published online today (June 10) in *JAMA Internal Medicine*.

Researchers analyzed data from 783 diabetic participants and found that hospitalization for severe hypoglycemia among the diabetic elderly in the study was associated with a doubled risk of developing dementia later. Similarly, [study participants](#) with dementia were twice as likely to experience a severe hypoglycemic event.

The study results suggest some patients risk entering a downward spiral in which hypoglycemia and [cognitive impairment](#) fuel one another, leading to worse health, said Kristine Yaffe, MD, senior author and principal investigator for the study, and a UCSF professor of psychiatry, neurology and epidemiology based at the San Francisco Veterans Affairs Medical Center.

"Older patients with diabetes may be especially vulnerable to a vicious cycle in which poor diabetes management may lead to [cognitive decline](#) and then to even worse [diabetes management](#)," she said.

The researchers analyzed hospital records of patients from Memphis and Pittsburgh, ages 70 to 79 at the time of enrollment, who participated in the federally funded Health, Aging and [Body Composition](#) (Health

ABC) study, begun in 1997. The UCSF results are based on an average of 12 years of follow-up study. Participants in the Health ABC study periodically underwent tests to measure cognitive function.

Nearly half of participants included in the newly published analysis were black, and the rest were white. None had dementia at the start of the study, and all either had diabetes at the beginning of the study or were diagnosed during the course of the study.

"Individuals with dementia or even those with milder forms of cognitive impairment may be less able to effectively manage complex treatment regimens for diabetes and less able to recognize the symptoms of hypoglycemia and to respond appropriately, increasing their risk of severe hypoglycemia," Yaffe said. "Physicians should take cognitive function into account in managing diabetes in elderly individuals."

Certain medications known to carry a higher risk for hypoglycemia—such as insulin secretagogues and certain sulfonylureas—may be inappropriate for older adults with dementia or who or at risk for cognitive impairment, according to Yaffe.

Previous studies in which researchers investigated [hypoglycemia](#) and cognitive function have had inconsistent findings. A strength of the current study is that individuals were tracked from baseline over a relatively long time, and the older age of participants may also have been a factor in the highly statistically significant outcome, Yaffe said.

More information: JAMA Intern Med. Published online June 10, 2013. doi:10.1001/jamainternmed.2013.6176

Provided by University of California, San Francisco

Citation: Alzheimer's and low blood sugar in diabetes may trigger a vicious cycle (2013, June 10)
retrieved 10 April 2024 from
<https://medicalxpress.com/news/2013-06-alzheimer-blood-sugar-diabetes-trigger.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.