

Food contaminants worsen metabolic problems in obese mice

June 27 2013

(Medical Xpress)—In order to get a better understanding of these effects, researchers from the Inserm cardiovascular, metabolism, diabetology and nutrition unit (U1060 " Laboratoire de recherche en cardiovasculaire, métabolisme, diabétologie et nutrition " Inserm/Inra/Université Lyon 1) introduced a "cocktail" of contaminants mixed with low doses of dioxin, PCB, bisphenol A and phtalates into the feeding of mice that had already been rendered obese by a high-fat diet.

The results show that metabolic changes occur in these <u>mice</u>, but that the effects differ depending on the gender. Females appeared to be more affected. Their obesity-induced glucose intolerance worsened and their estrogen pathway was altered.

These works have been published in the review Faseb Journal.

Obesity is a major public health problem because it is a risk factor in the development of <u>metabolic complications</u> (diabetes, cardiovascular diseases, etc.). It is a multi-factorial disorder. In addition to <u>genetic</u> <u>predispositions</u> and a life style that combines overeating with lack of exercise, there is a great volume of proof to suggest that contaminants, particularly in the food we eat, are responsible for the <u>obesity epidemic</u> and the resulting <u>metabolic changes</u>.

Researchers have put forward the hypothesis that contaminants in food could worsen certain metabolic problems already caused by eating an over-rich or a high-fat diet.



In this study, the researchers fed mice a high-fat diet (already a health risk), to which low doses of contaminants had been added. They were given this diet throughout their lives. Their mother had been nourished with this diet prior to their birth and during the gestation and <u>lactation</u> periods. Therefore, they suffered chronic exposure to this diet.

Two environmentally persistent contaminants (dioxin and PCB) and two non-persistent contaminants (phtalate and bisphenol A) were added to the high-fat (obesogenic) diet of the mice. The doses given were low, normally considered not to have any health impacts. These products were chosen because they are present in human food and because they are known to trigger endocrine disruption. In parallel, a control group of mice was fed with the same obesogenic diet, but without added contaminants.

The researchers then ran glucose tolerance and insulin sensitivity tests. They measured the livers for lipid accumulation and the expression of certain genes that play key roles in the metabolism of the adult mice.

The results show that the effects are highly dependent on the gender of the animal.

Male and female: different effects

In the females fed with a high-fat diet, the addition of contaminants worsened the glucose intolerance and altered the estrogen pathway. In males, it altered the cholesterol and lipid metabolism. There was no change in weight between the exposed mice and the unexposed mice.

The researchers pursued the hypothesis that there was a connection between the observed <u>glucose intolerance</u> and the alteration in the estrogen signaling in exposed females. It is well known that estrogens protect against metabolic disorders. In other words, these works suggest



that in obese females, exposure to food contaminants could lower the protection level that estrogens provide against metabolic disorders.

"With this study, we have succeeded in providing proof-of-concept that low doses of contaminants, even at levels normally considered to be without health impacts in humans, do in fact affect humans when subjected to chronic exposure, and when the contaminants are combined with a high-calorie <u>diet</u>" points out Brigitte Le Magueresse Battistoni.

More information: <u>dx.doi.org/</u>%2010.1096/fj.13-231670

Provided by Institut National de la Sante et de la Recherche Medicale

Citation: Food contaminants worsen metabolic problems in obese mice (2013, June 27) retrieved 3 May 2024 from https://medicalxpress.com/news/2013-06-food-contaminants-worsen-metabolic-problems.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.