

# Study finds greater potential benefit in overall survival for eribulin compared with capecitabine

June 12 2013

---

Subgroup analyses from a phase III clinical trial comparing a newer chemotherapy agent called eribulin mesylate, with capecitabine, a standard chemotherapy medication in women with previously treated metastatic breast cancer, showed increased benefit among women sharing certain traits. Specifically, these analyses demonstrated a greater potential benefit in certain subsets of patients with metastatic breast cancer. This analysis was presented by Peter A. Kaufman, M.D., during the 2013 ASCO Annual Meeting.

The specific patient populations who appeared to benefit from eribulin, in comparison to [capecitabine](#), are as follows:

- Patients with more than two organs involved with metastatic [breast cancer](#)
- Patients who had not received chemotherapy for six months or longer
- Patients who had received [anthracycline](#) and/or a taxane therapies in the metastatic setting

Previous pre-specified exploratory analysis of overall survival and progression-free survival showed women with triple-negative, ER-negative, HER2-negative also had a greater relative benefit in overall

survival with eribulin over capecitabine.

"These exploratory analyses suggest that other patient subgroups may benefit from eribulin and further studies are warranted," said Peter A. Kaufman, MD, associate professor of medicine at the Geisel School of Medicine at Dartmouth, and oncologist at Dartmouth-Hitchcock and Norris Cotton Cancer Center in Lebanon, N.H.

In 2010, the FDA approved eribulin for the treatment of patients with metastatic breast cancer who had previously received an anthracycline and a taxane and at least two cytotoxic chemotherapy [treatment regimens](#) for metastatic breast cancer. The FDA granted approval based on data showing a statistically significant improvement in overall survival compared with current treatments.

Kaufman and colleagues are still compiling data from the quality-of-life analysis, which according to Kaufman, will help guide their next steps in further studying eribulin in this patient population.

Provided by Dartmouth-Hitchcock Medical Center

Citation: Study finds greater potential benefit in overall survival for eribulin compared with capecitabine (2013, June 12) retrieved 13 March 2024 from <https://medicalxpress.com/news/2013-06-greater-potential-benefit-survival-eribulin.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--