

## Hispanic and black kids less likely to use medication to control asthma

June 28 2013, by Katherine Kahn

---



(Medical Xpress)—Black and Hispanic children with asthma are less likely than White children to use long-term asthma control medications, finds a new study in *Health Services Research*.

Eric Sarpong, PhD, the study's lead author commented, "Several studies continue to find that minority children are less likely to use recommended controller medications and more likely to use relievers to manage the disease."

Reliever medications are quick-acting drugs used to manage acute symptoms of asthma. The National Asthma Education and Prevention Program recommends that long-term controller medications—medications taken daily to reduce [airway inflammation](#)

and prevent onset of [asthma symptoms](#)—should be used to achieve and maintain control of [persistent asthma](#).

Sarpong and Edward Miller, Ph.D., researchers at the federal Agency for Health Research and Quality, wanted to research the contributing factors for this disparity in medication use.

The study used nationally representative data from the 2005-2008 Medical Expenditure Panel Survey, the 2000 Decennial Census, and the 2004-2007 National Health Interview Survey and included data on both privately and publicly insured children.

Living in the western United States and having non-native parents was linked with the majority of the difference in medication use between Hispanic and White children. Lower family income and a lower likelihood of having parents who are married was also associated with the difference in controller use for both Black and Hispanic children.

However, much of the difference in controller use by Black children compared to White children was largely unexplained. This finding was observed even when taking insurance status into account. "This may point to biases in the health system, maybe in prescribing patterns, or in the quality of care," Sarpong says.

Glenn Flores, M.D., professor of pediatrics at the University of Texas Southwestern in Dallas, considers healthcare provider and healthcare system bias to be a prominent factor in sub-optimal asthma care. "Too often we're blaming the kids or the families when, in fact, it may be we as healthcare providers or the healthcare system that's responsible for not just this racial and ethnic disparity, but many racial and ethnic disparities," he says.

Flores also considers inadequate patient and family education, healthcare

provider failure to schedule follow-up care, and failure to provide prescriptions for medication refills to be key areas for improvement. He also emphasizes the importance of access to specialty care, which has been shown to improve outcomes in [asthma](#). "We and others have shown that minority kids in general—no matter what condition you're looking at—have lower access to specialty care," he says.

**More information:** Sarpong, E. and Miller, G. Racial and ethnic differences in childhood asthma treatment in the United States, *Health Services Research*. (2013).

Provided by Health Behavior News Service

Citation: Hispanic and black kids less likely to use medication to control asthma (2013, June 28) retrieved 19 April 2024 from <https://medicalxpress.com/news/2013-06-hispanic-black-kids-medication-asthma.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.