

Study assesses impact of rheumatoid arthritis on joint replacement surgery outcomes

June 12 2013

Two new studies by researchers at Hospital for Special Surgery have shed light on joint replacement outcomes in patients with rheumatoid arthritis (RA). One study overturns the common belief that RA patients have worse outcomes after a total knee replacement (TKR) than patients who undergo the operation for osteoarthritis. The other study demonstrates that RA patients who undergo a total hip replacement were as likely to have significant improvements in function and pain as patients with osteoarthritis (OA), even though they did not do as well.

The news will be reported at the European League Against Rheumatism (EULAR) annual Congress of Rheumatology, to be held June 12-15, in Madrid, Spain.

The studies were made possible by the HSS Total Joint Replacement Registry. Started in 2007, this prospective registry includes a gold mine of information including data on all [patients](#) who seek care at HSS for knee and [hip replacement surgery](#). "Other hospitals have registries, but what we have is an incredible volume of information on patients that allows us to do interesting studies, because there are so many patients having hip and [knee replacement](#) surgeries at our hospital," said Susan Goodman, M.D., lead author of both studies and a rheumatologist at Hospital for Special Surgery in New York City.

Historically, RA patients have had worse outcomes after joint

replacement surgeries, however, whether this was caused by poorly controlled disease or disease treatment was unclear. Starting in the 1980s, effective disease-modifying drugs became available to treat patients with RA. In the late 1990's, [etanercept](#), [infliximab](#), and other biologic medication came on the market. Today, at HSS over 70% of patients are on disease modifying drugs and over 50% are on biologics.

In the two new studies, investigators at Special Surgery set out to examine whether outcomes remained worse for RA patients in a cohort with a high prevalence of potent disease-modifying drugs and [biologic agents](#). Patient pain and function was assessed prior to surgery and two years after surgery using the Western Ontario and McMaster Universities Arthritis Index (WOMAC). This tool measures pain, stiffness, and functional limitation. Knee Replacement

In one study, investigators used the HSS Total [Joint Replacement](#) registry to identify 178 [rheumatoid arthritis](#) patients and 5,206 osteoarthritis patients who underwent a TKR. Patients with RA were sicker—while 72% of osteoarthritis patients had no comorbidities, only 34% of rheumatoid arthritis patients fell into this category.

Although TKR patients with RA had worse pain and function prior to surgery compared to controls, these differences disappeared after surgery. "In RA patients, their preoperative scores were significantly worse than the osteoarthritis controls, but in fact our knee replacement patients caught up," said Dr. Goodman. Patients in both groups had similar satisfaction rates.

Investigators also identified 32 RA patients and 342 OA patients who underwent a TKR revision. In this analysis, they discovered that OA and RA patients had similar pain and function pre-operatively, but patients with rheumatoid arthritis actually had significantly less pain, better function, and were much more satisfied at two years. "Our RA patients

undergoing knee replacement revisions did better than our osteoarthritis patients," said Dr. Goodman. Hip Replacement

In a second study, HSS investigators compared outcomes of 202 RA patients and 5,810 OA patients who underwent hip replacement. Again, RA patients were sicker—while 80% of osteoarthritis patients had no comorbidities, only 34% of rheumatoid arthritis patients had the luxury of falling into this category.

"When we looked at function using the WOMAC scale, function was significantly worse in the rheumatoid arthritis patients prior to surgery, but interestingly, when we looked at how they did two years down the road, they were as likely to have an improvement in function and pain," said Dr. Goodman. Almost 100% (RA, 96% and OA, 95%) had a ten point change in score—a ten point change is considered clinically significant. Rheumatoid arthritis patients, however, had worse outcomes. At two years, patients with RA were more likely to have a poor function WOMAC score of 60 or less (18% vs. 4%) and a poor WOMAC pain score of 60 or less (12% vs. 3%). Patients who expected to do well with the surgery had a significantly decreased risk of poor pain and function.

"The rheumatoid arthritis patients have a great response in terms of hip replacement and their hip function improved.

Nonetheless, they were a group that did not do as well," said Dr. Goodman.

Similar findings were identified for patients undergoing hip replacement revision (58 RA patients and 445 OA patients).

"Many RA patients have a worsening or flare of their disease six weeks after surgery," said Dr. Goodman. "It may be that those patients aren't able to do their physical therapy because they are not feeling well, and

maybe that contributes to poor outcomes down the road."

Clearly, there is a difference in knee replacement versus hip replacement in patients with RA, but it is unclear exactly what the difference is.

"Perhaps the RA patients undergoing [hip replacement surgery](#) are delaying their [surgery](#) too long. Maybe if we intervene sooner, we would do better functionally," said Dr. Goodman. The HSS investigators plan to conduct further research to identify possible differences between the two groups.

More information: This study will be presented on Saturday, June 15, at 10:15 a.m. in a poster session.

Provided by Hospital for Special Surgery

Citation: Study assesses impact of rheumatoid arthritis on joint replacement surgery outcomes (2013, June 12) retrieved 28 April 2024 from <https://medicalxpress.com/news/2013-06-impact-rheumatoid-arthritis-joint-surgery.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--