

Risk of kidney disease doubled with use of fluoroquinolone antibiotics

June 3 2013

The risk of acute kidney disease is doubled for people taking oral fluoroquinolone antibiotics, according to a study published in *CMAJ* (*Canadian Medical Association Journal*).

Fluoroquinolones, including ciprofloxacin, levofloxacin and moxifloxacin, are common broad-spectrum antibiotics most often used to treat respiratory and urogenital infections. Case reports have indicated acute kidney injury with use, and prescription labels carry a warning of [kidney failure](#). However, when oral fluoroquinolones are prescribed in clinical practice, kidney injury is usually not considered.

Researchers from the US and Canada undertook a study to determine the risk of acute kidney injury with the use of oral fluoroquinolones for adult men 40

More information: www.cmaj.ca/lookup/doi/10.1503/cmaj.121730

Provided by Canadian Medical Association Journal

Citation: Risk of kidney disease doubled with use of fluoroquinolone antibiotics (2013, June 3) retrieved 25 April 2024 from <https://medicalxpress.com/news/2013-06-kidney-disease-fluoroquinolone-antibiotics.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.