

# One in four stroke patients suffer PTSD

June 19 2013

---

One in four people who survive a stroke or transient ischemic attack (TIA) suffer from symptoms of post-traumatic stress disorder (PTSD) within the first year post-event, and one in nine experience chronic PTSD more than a year later. The data suggest that each year nearly 300,000 stroke/TIA survivors will develop PTSD symptoms as a result of their health scare. The study, led by Columbia University Medical Center researchers, was published today in the online edition of *PLOS ONE*.

"This work builds on recent findings of ours that PTSD is common among heart attack survivors and that it contributes to a doubled risk of a future cardiac event or of dying within one to three years. Our current results show that PTSD in [stroke](#) and TIA survivors may increase their risk for [recurrent stroke](#) and other cardiovascular events," said first author Donald Edmondson, PhD, MPH, assistant professor of behavioral medicine (Center for Behavioral Cardiovascular Health) at CUMC. "Given that each event is life-threatening and that strokes/TIAs add hundreds of millions of dollars to annual [health expenditures](#), these findings are important to both the long-term survival and health costs of these patient populations."

"PTSD is not just a disorder of combat veterans and sexual assault survivors, but strongly affects survivors of stroke and other potentially traumatic acute cardiovascular events as well," said Ian M. Kronish, MD, MPH, assistant professor of medicine (Center for Behavioral Cardiovascular Health) and the study's senior author. "Surviving a life-threatening health scare can have a debilitating psychological impact,

and [health care providers](#) should make it a priority to screen for [symptoms of depression](#), anxiety, and PTSD among these patient populations."

Stroke is the fourth-leading cause of death and the top cause of disability in the United States. According to data from the American Stroke Association, nearly 795,000 Americans each year suffer a new or recurrent stroke, and up to an additional 500,000 suffer a TIA.

PTSD is an anxiety disorder initiated by exposure to a traumatic event. Common symptoms include nightmares, avoidance of reminders of the event, and elevated heart rate and blood pressure. [Chronic PTSD](#) is a duration of these symptoms for three months or longer (as defined by the DSM-IV).

Since only a few studies have assessed PTSD due to stroke, Drs. Edmondson and Kronish and their colleagues performed the first meta-analysis of clinical studies of stroke- or TIA-induced PTSD. The nine studies in the meta-analysis included a total of 1,138 stroke or TIA survivors.

The study found that 23 percent, or roughly one in four, of the patients developed PTSD symptoms within the first year after their stroke or TIA, with 11 percent, or roughly one in nine, experiencing chronic PTSD more than a year later.

"PTSD and other psychological disorders in stroke and TIA patients appear to be an under-recognized and undertreated problem," said Dr. Kronish.

"Fortunately, there are good treatments for PTSD," said Dr. Edmondson. "But first, physicians and patients have to be aware that this is a problem. Family members can also help. We know that social support is

a good protective factor against PTSD due to any type of traumatic event."

"The next step is further research to assess whether mental health treatment can reduce stroke- and TIA-induced PTSD symptoms and help these patients regain a feeling of normalcy and calm as soon as possible after their health scare," said Dr. Edmondson.

**More information:** "Prevalence of PTSD in Survivors of Stroke and Transient Ischemic Attack: A Meta-analytic Review."  
[dx.plos.org/10.1371/journal.pone.0066435](https://doi.org/10.1371/journal.pone.0066435)

Provided by Columbia University Medical Center

Citation: One in four stroke patients suffer PTSD (2013, June 19) retrieved 27 April 2024 from <https://medicalxpress.com/news/2013-06-patients-ptsd.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.