

# A prenatal trigger for postnatal obesity

June 24 2013

---

During pregnancy, the health of the mother and the intrauterine environment can have dramatic and lasting effects on the child. Intrahepatic cholestasis of pregnancy (ICP) is a liver disease that affects 0.5-2% of pregnant women and is characterized by increased bile acid levels in the maternal serum.

In this issue of the *Journal of Clinical Investigation*, Catherine Williamson and colleagues at Imperial College London studied the long term impact of ICP in a cohort of Finnish families.

They found that as teenagers, individuals born to women with ICP had altered metabolic profiles and increased BMI. To further understand this effect, Williams and colleagues developed a mouse model of ICP and found that offspring of ICP mothers were more susceptible to metabolic disease and diet-induced obesity.

In the companion commentary, Susan Murphy of Duke University points out that the mouse model of ICP may also be useful in identifying other factors that predispose individuals to metabolic syndrome.

**More information:** Cholestatic pregnancy programmes metabolic disease in the offspring, *J Clin Invest.* 2013;123(7):3172–3181.

[doi:10.1172/JCI68927](https://doi.org/10.1172/JCI68927)

Provided by Journal of Clinical Investigation

Citation: A prenatal trigger for postnatal obesity (2013, June 24) retrieved 19 April 2024 from <https://medicalxpress.com/news/2013-06-prenatal-trigger-postnatal-obesity.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.