

# Psychiatric disorders linked to a protein involved in the formation of long-term memories

June 17 2013, by Ken Kingery

---

Researchers have discovered a pathway by which the brain controls a molecule critical to forming long-term memories and connected with bipolar disorder and schizophrenia.

The discovery was made by a team of scientists led by Alexei Morozov, an assistant professor at the Virginia Tech Carilion Research Institute.

The mechanism – a protein called Rap1 – controls L-type [calcium channels](#), which participate in the formation of long-term memories. Previous studies have also linked alterations in these [ion channels](#) to certain psychiatric disorders. The discovery of the channels' regulation by Rap1 could help scientists understand the physiological genesis of bipolar disorder and [schizophrenia](#).

"People with [genetic mutations](#) affecting L-type calcium channels have higher rates of bipolar disorder and schizophrenia," said Morozov. "This suggests that there might be a relationship between the activation of L-type calcium channels and these psychiatric disorders. Understanding how these ion channels are controlled is the first step to determining how their functioning or malfunctioning affects mental health."

A single neuron in the brain can have thousands of synapses, each of which can grow, strengthen, weaken, and change structurally in response to learning new information. [Electric signals](#) traveling from neuron to

neuron jump across these synapses through chemical neurotransmitters. The release of these chemicals is caused by the flow of electrically charged atoms through a particular subset of ion channels known as voltage-gated calcium channels.

Previous studies have shown that blocking these ion channels inhibits the formation of long-term memories. Although it was known that L-type calcium channels are activated in response to learning, how they are controlled was a mystery.

In the experiment, Morozov and colleagues knocked out the gene responsible for coding the enzyme Rap1, which he suspected played a role in activating L-type calcium channels. The researchers then used live imaging techniques to monitor the release of neurotransmitters and electron microscopy to visualize L-type channels at synapses. They discovered that, without Rap1, the L-type calcium channels were more active and more abundant at [synapses](#) all the time, increasing the release of neurotransmitters. The results showed that Rap1 is responsible for suppressing L-type calcium channels, allowing them to activate only at the proper moments, possibly during long-term memory formation.

"Our next step is to determine whether this new signaling pathway is altered in cases of mental disease," said Morozov. "If so, it could help us gain a better understanding of the molecular underpinnings of channel-related [psychiatric disorders](#), such as bipolar disorder and schizophrenia. Such knowledge would go a long way toward developing new therapeutic methods."

**More information:** The discovery appeared in *The Journal of Neuroscience* in the study "Rap1 Signaling Prevents L-Type Calcium Channel-Dependent Neurotransmitter Release," by Jaichandar Subramanian: [www.jneurosci.org/content/33/17/7245.abstract](http://www.jneurosci.org/content/33/17/7245.abstract)

Provided by Virginia Tech

Citation: Psychiatric disorders linked to a protein involved in the formation of long-term memories (2013, June 17) retrieved 1 May 2024 from <https://medicalxpress.com/news/2013-06-psychiatric-disorders-linked-protein-involved.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.