

Study shows radiotherapy and surgery provide regional control for breast cancer patients

June 3 2013

Final analysis of the EORTC 10981-22023 AMAROS (After Mapping of the Axilla: Radiotherapy Or Surgery?) trial has shown that both axillary lymph node dissection and axillary radiotherapy provide excellent regional control for breast cancer patients with a positive sentinel node biopsy. The AMAROS trial also found that axillary radiotherapy reduces the risk of short term and long-term lymphoedema as compared to axillary lymph node dissection.

Prof. Emiel J. Rutgers of The Netherlands Cancer Institute-Antoni Van Leeuwenhoekziekenhuis in Amsterdam, the EORTC Breast Cancer Group, and principal investigator of this study says, "[Sentinel node biopsy](#) has become the standard procedure for assessing axillary lymph node status for cN0 [breast cancer](#) patients. At the start of this trial the question was whether axillary lymph node dissection should remain the standard treatment in case of a tumor positive sentinel node."

From 2001 to 2010, 4806 patients who were diagnosed with [invasive breast cancer](#) were enrolled in this phase III non-[inferiority](#) trial, and randomized for their axillary treatment in case the sentinel node [biopsy](#) was positive. Of these, 3131 patients had negative biopsies and 250 patients had a different sentinel node outcome. Thus, out of 1425 patients with positive node, axillary lymph node dissection was to be used in 744 patients, and axillary [radiotherapy](#) in 681 patients.

Dr. Geertjan van Tienhoven of the Academisch Medisch Centrum - Universiteit van Amsterdam and principal investigator on behalf of the EORTC [Radiation Oncology](#) Group says, "Although axillary lymph node dissection provides excellent regional control, it was known to cause harmful side effects. Axillary radiotherapy instead of axillary lymph node dissection was hypothesized to provide comparable regional control and perhaps fewer side effects in this particular group of patients."

The primary endpoint of this EORTC trial was 5-year axillary recurrence rate. At a median follow up of 6.1 years, this rate was 0.54% (4 out of 744 patients) after axillary lymph node dissection and 1.03% (7 out of 681 patients) after axillary radiotherapy. Due to the unexpectedly low number of events, the non-inferiority test was underpowered. The axillary recurrence rate for patients with a negative sentinel node biopsy was 0.8% (25 out of 3131 patients).

Secondary endpoints were overall survival, disease-free survival, quality of life, shoulder movement and lymphoedema at one and five years. No significant differences in overall or disease free survival were observed between the two treatment arms. Five year estimates for overall survival were 93.3% for the arm receiving axillary lymph node dissection and 92.5% for patients in the axillary radiotherapy arm ($p=0.34$), and for disease-free survival the five year estimates were 86.9% for axillary lymph node dissection and 82.6% for axillary radiotherapy ($p=0.18$).

Lymphoedema, a swelling of tissue (typically occurring in the arm), occurred more often following axillary lymph node dissection, 40% at one year and 28% at five years, than after axillary radiotherapy, 22% at one year and 14% at five years (p

Citation: Study shows radiotherapy and surgery provide regional control for breast cancer patients (2013, June 3) retrieved 25 April 2024 from

<https://medicalxpress.com/news/2013-06-radiotherapy-surgery-regional-breast-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.