

Study examines characteristics, features of West Nile virus outbreaks

July 16 2013

An analysis of West Nile virus epidemics in Dallas County in 2012 and previous years finds that the epidemics begin early, after unusually warm winters; are often in similar geographical locations; and are predicted by the mosquito vector index (an estimate of the average number of West Nile virus-infected mosquitoes collected per trap-night), information that may help prevent future outbreaks of West Nile virus-associated illness, according to a study in the July 17 issue of *JAMA*.

"After declining over the prior 5 years, mosquito-borne West Nile virus infection resurged in 2012 throughout the United States, most substantially in Dallas County, Texas. Dallas has been a known focus of mosquito-borne encephalitis since 1966, when a large epidemic of St. Louis encephalitis (SLE) occurred there, necessitating aerial spraying of insecticide for control," according to background information in the article. "With the introduction of West Nile virus into New York City in 1999 and its subsequent spread across the country, West Nile virus appears to have displaced SLE virus. Dallas recognized its initial cases of West Nile virus encephalitis in 2002 and its first sizeable outbreak in 2006, followed by 5 years of low West Nile virus activity. In the 2012 nationwide West Nile virus resurgence, Dallas County experienced the most West Nile virus infections of any U.S. urban area, requiring intensified ground and <u>aerial spraying</u> of insecticides."

Wendy M. Chung, M.D., S.M., of Dallas County Health and Human Services, Dallas, and colleagues conducted a study to examine the features associated with the West Nile virus epidemics and to identify



surveillance and <u>control measures</u> for minimizing future outbreaks. The researchers analyzed <u>surveillance data</u> from Dallas County (population, 2.4 million), which included the numbers of residents diagnosed with West Nile virus infection between May 30, 2012 and December 3, 2012; mosquito trap results; <u>weather data</u>; and syndromic (pertaining to symptoms and syndromes) surveillance from area emergency departments.

From May 30 through December 3, 2012, patients (n = 1,162) with any West Nile virus-positive test result were reported to the health department; 615 met laboratory case criteria, and 398 cases of West Nile virus illness with 19 deaths were confirmed by clinical review in residents of Dallas County. The outbreak included 173 patients with West Nile neuroinvasive disease (WNND) and 225 with West Nile fever, and 17 West Nile virus-positive blood donors. Regarding patients with WNND, 96 percent were hospitalized; 35 percent required intensive care; 18 percent required assisted ventilation; and the casefatality rate was 10 percent. The overall WNND incidence rate in Dallas County was 7.30 per 100,000 residents in 2012, compared with 2.91 in 2006.

The first West Nile virus-positive mosquito pool of 2012 was detected in late May, earlier than in typical seasons. Symptoms of the first 19 cases of WNND in 2012 began in June, a month earlier than in most prior seasons; thereafter, the number of new cases escalated rapidly. Sequential increases in the weekly vector index early in the 2012 season significantly predicted the number of patients with onset of symptoms of WNND in the subsequent l to 2 weeks.

The 2012 epidemic year was distinguished from the preceding 10 years by the mildest winter, as indicated by absence of hard winter freezes, the most degree-days above daily normal temperature during the winter and spring and other features. During the 11 years since West Nile virus was



first identified in Dallas, the researchers found that the annual prevalence of WNND was inversely associated with the number of days with low temperatures below 28°F in December through February.

"Although initially widely distributed, WNND cases soon clustered in neighborhoods with high housing density in the north central area of the county, reflecting higher vector indices and following geospatial patterns of West Nile virus in prior years," the authors write.

Aerial insecticide spraying was not associated with increases in emergency department visits for respiratory symptoms or skin rash.

"This report identifies several distinguishing features of a large urban West Nile virus outbreak that may assist future prevention and control efforts for vector-borne infections," the authors write. "Consideration of weather patterns and historical geographical hot spots and acting on the vector index may help prevent West Nile virus-associated illness."

"Periodic flares of West Nile virus, as occurred in 2012, certainly will recur," writes Stephen M. Ostroff, M.D., formerly of the Centers for Disease Control and Prevention, Atlanta, and the Pennsylvania Department of Health, Harrisburg, in an accompanying editorial.

"Where future outbreaks of the virus will occur and how intense they will be is difficult to predict, especially in light of declining surveillance efforts and vector monitoring programs. Unusually <u>warm winters</u>, as occurred in Dallas during 2011-2012, are becoming more common and will favor additional West Nile virus events like the one described by Chung et al. Changing weather patterns raise the possibility of expanded zones of risk and longer transmission seasons. The tragic consequences of the Dallas West Nile virus epidemic must not be forgotten, for they serve as a cogent reminder of the need to sustain vector monitoring and prevention programs in all communities."



More information: *JAMA*. 2013;310(3):297-307 *JAMA*. 2013;310(3):267-268

Provided by The JAMA Network Journals

Citation: Study examines characteristics, features of West Nile virus outbreaks (2013, July 16) retrieved 5 May 2024 from https://medicalxpress.com/news/2013-07-characteristics-features-west-nile-virus.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.