


Decoding mechanisms of cell orientation in the brain

July 31 2013



The upper two pictures show NG2-expressing OPC in healthy nerve tissue. In comparison, the lower pictures show the altered morphology and orientation of cells in damaged nerve tissue. Credit: Dominik Sakry

When the central nervous system is injured, oligodendrocyte precursor cells (OPC) migrate to the lesion and synthesize new myelin sheaths on

demyelinated axons. Scientists at the Institute of Molecular Cell Biology at Johannes Gutenberg University Mainz (JGU) have now discovered that a distinct protein regulates the direction and movement of OPC toward the wound. The transmembrane protein NG2, which is expressed at the surface of OPCs and down-regulated as they mature to myelinating oligodendrocytes, plays an important role in the reaction of OPC to wounding. The results of this study have recently been published in the renowned *Journal of Neuroscience*.

The [myelin sheath](#) functions to electrically isolate axons of many [nerve fibers](#) and is synthesized by oligodendrocytes which mature from the OPC. In the case of injury, neural cells send out signaling molecules which attract the OPC. The NG2 protein helps OPCs to react to some of these and move in a directed and orientated fashion. "We were able to prove in cell biological experiments that NG2 orientates OPC toward the lesion and ensures targeted OPC migration toward the wound through the regulation of cell polarity", explained Dr. Fabien Binamé, lead author of the study. Supported by funding of the German Research Foundation (DFG), Dr. Fabien Binamé is currently carrying out his research at the Institute of Molecular Cell Biology headed by Professor Jacqueline Trotter.

"The function and mode of operation of NG2 is not yet fully understood", added co-author Dominik Sakry, who was also involved in the study. "But it looks as if the NG2-associated regulatory mechanism becomes apparent only in cases of injury of the nervous system."

Diseases such as Multiple Sclerosis or brain tumors go hand in hand with damage of nerve tissue. "The results of our study on NG2-mediated basic mechanisms of cell orientation and migration could aid in understanding the repair of damaged demyelinated tissue, or be important for treatment of highly active migratory [brain tumors](#) which often express high levels of NG2", said Professor Jacqueline Trotter,

head of the JGU Institute of Molecular Cell Biology.

More information: F. Binamé, D. Sakry, L. Dimou, V. Jolivel, J. Trotter, NG2 Regulates Directional Migration of Oligodendrocyte Precursor Cells via Rho GTPases and Polarity Complex Proteins, *Journal of Neuroscience*, 26 June 2013
[DOI:10.1523/JNEUROSCI.5010-12.2013](https://doi.org/10.1523/JNEUROSCI.5010-12.2013)

Provided by Universitaet Mainz

Citation: Decoding mechanisms of cell orientation in the brain (2013, July 31) retrieved 26 April 2024 from <https://medicalxpress.com/news/2013-07-decoding-mechanisms-cell-brain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.