

Stress fracture risks may be modifiable

July 12 2013

Programs to improve movement patterns may help prevent stress fractures in athletes and military personnel, say researchers presenting their work today at the American Orthopaedic Society for Sports Medicine's Annual Meeting in Chicago, IL.

"Several potentially modifiable muscular and biomechanical factors may be playing a part in the increased rates of [stress fractures](#) in athletes and military personnel," said Kenneth L. Cameron, PhD, MPH, ATC, lead author and Director of Orthopaedic Research at Keller Army Hospital, West Point, NY. "It's possible that injury prevention programs targeted to address these movement patterns can help reduce stress [fracture risk](#)."

Cameron and his colleagues analyzed data from the JUMP-ACL cohort, which is an existing study of military cadets, detailing [motion analysis](#) during a jump landing task at the initiation of each participant's military career. They studied 1,843 individuals from the class years of 2009-2012, with 94 subjects sustaining a lower extremity stress fracture during the follow-up period.

The incidence rate for stress fracture injuries among females was nearly three times greater when compared to males. Knee rotation and abduction angles when landing were both associated with the rates of lower-extremity stress fractures, as were reduced knee and hip flexion angles, and increased vertical and medial ground reaction forces.

"Lower extremity movement patterns and strength have previously been associated with stress fractures and [overuse injuries](#); however, our study

is one of the first to identify dynamic knee rotation and frontal plane angles as important prospective risk factors for lower extremity stress fractures. We hope that by better understanding the [movement patterns](#) associated with lower extremity stress fracture injury risk we can help create programs for prevention," said Cameron. The study was supported by research grants from the Congressionally Directed Medical Research Program's (CDMRP) Peer Reviewed Medical Research Program and the National Institute of Arthritis, Musculoskeletal, and Skin Diseases (NIAMS).

Provided by American Orthopaedic Society for Sports Medicine

Citation: Stress fracture risks may be modifiable (2013, July 12) retrieved 18 April 2024 from <https://medicalxpress.com/news/2013-07-stress-fracture.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--