

Vascular complications of fungal meningitis after contaminated spinal injections

July 22 2013

A case series by researchers at Vanderbilt University, Nashville, Tenn., examined three patients with ischemic stroke who later received a diagnosis of fungal meningitis attributed to epidural injections of contaminated methylprednisolone for low back pain.

The recent identification of injections of contaminated methylprednisolone acetate has highlighted the different clinical presentations of [fungal meningitis](#), which can have an [incubation period](#) of one to four weeks between the last spinal injection and when a patient seeks medical care.

"Fungal meningitis due to injections of contaminated methylprednisolone acetate can present with vascular sequelae in immunocompetent individuals. This is particularly germane to neurologists because better recognition of clinical characteristics of patients with fungal meningitis and ischemic stroke will provide more timely and efficient care," the paper concludes.

More information: *JAMA Neurol.* Published online July 22, 2013.
[doi:10.1001/jamaneurol.2013.3586](https://doi.org/10.1001/jamaneurol.2013.3586)

Provided by The JAMA Network Journals

Citation: Vascular complications of fungal meningitis after contaminated spinal injections (2013,

July 22) retrieved 27 April 2024 from <https://medicalxpress.com/news/2013-07-vascular-complications-fungal-meningitis-contaminated.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.