

# Women at risk of developing postpartum psychosis need close monitoring, says new review

July 11 2013

---

There are clear risk factors for postpartum psychosis that all women should be asked about antenatally to ensure early recognition and prompt treatment of the condition, says a new review published today (12 July) in *The Obstetrician & Gynaecologist (TOG)*.

Postpartum psychosis is a severe mental illness with a dramatic onset shortly after childbirth, affecting approximately 1-2 in 1000 deliveries. However, the review notes that the true incidence may be higher.

Common symptoms include; mania, severe depression, delusions and hallucinations, confusion, bewilderment or perplexity, all of which increase the risk for both mother and child.

The review notes that there is consistent evidence of a specific relationship between postpartum psychosis and bipolar disorder. Women with bipolar disorder have at least a 1 in 4 risk of suffering postpartum psychosis. Genetics are also a factor and [women](#) with [bipolar disorder](#) and a personal or family history of postpartum psychosis are at particularly high risk with greater than 1 in 2 deliveries affected by postpartum psychosis.

However, half of women who develop postpartum psychosis have no family history or previous [risk factors](#) that put them in a high risk group of suffering from the condition.

The review emphasises the need for close contact and review from a multidisciplinary team during the perinatal period for at least three months following delivery, even if the woman is well and recommends a written plan covering pregnancy and the postnatal period which should be discussed with the woman and her family.

Dr Ian Jones, Reader in Perinatal Psychiatry, Cardiff University and co-author of the review said:

"Women at high risk of postpartum psychosis need very careful care before conception, throughout pregnancy and during the postpartum period, including pre-conception counselling and close monitoring and psychiatric assessment after childbirth.

"Postpartum psychosis is a true psychiatric emergency and it is vital that is recognised early and treated immediately. Admission to hospital is usually necessary and women should ideally be offered a specialist mother and baby unit where the best treatment options can be established."

Jason Waugh, *TOG*'s Editor-in-chief said:

"This review emphasises the importance of women at [high risk](#) of postpartum psychosis as well as the early recognition and prompt treatment of women who develop the condition.

"This paper also underlines that half of women who experience postpartum psychosis have no previous risk factors. It is therefore vital that all women are made aware of the condition and its signs and symptoms."

**More information:** Di Florio A, Smith S, Jones I. Postpartum psychosis. *The Obstetrician & Gynaecologist* 2013;

[dx.doi.org/10.1111/tog.2041](https://doi.org/10.1111/tog.2041)

Provided by Wiley

Citation: Women at risk of developing postpartum psychosis need close monitoring, says new review (2013, July 11) retrieved 23 April 2024 from

<https://medicalxpress.com/news/2013-07-women-postpartum-psychosis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.