

OK to continue aspirin therapy up to pancreatic surgery

August 1 2013



Patients on aspirin therapy do not have higher rates of perioperative bleeding, transfusion, or major procedure-related complications following elective pancreatic surgery, according to research published online July 26 in *Surgery*.

(HealthDay)—Patients on aspirin therapy do not have higher rates of perioperative bleeding, transfusion, or major procedure-related complications following elective pancreatic surgery, according to research published online July 26 in *Surgery*.

Andrea M. Wolf, M.D., of Spectrum Health in Grand Rapids, Mich., and colleagues conducted a retrospective, observational study using data from 1,017 patients who underwent pancreatic resection. Of these, 289 (28.4 percent) were continued on prescribed [aspirin therapy](#) through the morning of the surgery.

The researchers found that patients in the aspirin cohort were older than

those in the non-aspirin cohort (median age, 69 versus 62 years; P

"Despite many surgeons' fears that preoperative aspirin use increases bleeding, this study (among others in non-[pancreatic surgery](#)) shows that this [apprehension](#) appears unfounded," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

[Health News](#) Copyright © 2013 [HealthDay](#). All rights reserved.

Citation: OK to continue aspirin therapy up to pancreatic surgery (2013, August 1) retrieved 3 May 2024 from <https://medicalxpress.com/news/2013-08-aspirin-therapy-pancreatic-surgery.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
