

At-home care cuts depression in older African-Americans

August 20 2013



A home-based intervention delivered by social workers reduces symptoms and improves quality of life in older African-Americans with depressive symptoms, according to research published in the Aug. 20 issue of the *Annals of Internal Medicine*.

(HealthDay)—A home-based intervention delivered by social workers reduces symptoms and improves quality of life in older African-Americans with depressive symptoms, according to research published in the Aug. 20 issue of the *Annals of Internal Medicine*.

Laura N. Gitlin, Ph.D., of Johns Hopkins University in Baltimore, and colleagues randomly assigned 208 African-Americans aged 55 years and older with depressive symptoms to either a multicomponent, home-based care management [intervention group](#) (106 patients) or a wait-list control group (102 patients).

The researchers found that, among the 89 participants who completed

four months of the intervention program (compared with 93 participants in the control group) significant differences were observed in reduced severity of depression, increased knowledge of depression, improved quality of life, greater behavioral activation, reduced anxiety, and improved function. At four months, more participants in the intervention group entered remission (43.8 percent) than those in the control group (26.9 percent).

"African-Americans often have limited [mental health care](#) options," the authors write. "A home-based intervention delivered by social workers could reduce depressive symptoms and enhance quality of life in some older African-Americans."

More information: [Full Text \(subscription or payment may be required\)](#)

Copyright © 2013 [HealthDay](#). All rights reserved.

Citation: At-home care cuts depression in older African-Americans (2013, August 20) retrieved 13 March 2024 from <https://medicalxpress.com/news/2013-08-at-home-depression-older-african-americans.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
