

Autistic children can outgrow difficulty understanding visual cues and sounds

August 28 2013

Scientists at Albert Einstein College of Medicine of Yeshiva University have shown that high-functioning autism spectrum disorder (ASD) children appear to outgrow a critical social communication disability. Younger children with ASD have trouble integrating the auditory and visual cues associated with speech, but the researchers found that the problem clears up in adolescence. The study was published today in the online edition of the journal *Cerebral Cortex*.

"This is an extremely hopeful finding," said lead author John Foxe, Ph.D., professor of pediatrics and in the Dominick P. Purpura Department of Neuroscience, as well as director of research of the Children's Evaluation and Rehabilitation Center at Einstein. "It suggests that the neurophysiological circuits for speech in these children aren't fundamentally broken and that we might be able to do something to help them recover sooner."

According to Dr. Foxe, the ability to integrate "heard" and "seen" speech signals is crucial to effective communication. "Children who don't appropriately develop this capacity have trouble navigating educational and <u>social settings</u>," he said.

In a previous study, Dr. Foxe and his colleagues demonstrated that children with ASD integrate multisensory information such as sound, touch and vision differently from typically developing children. Among typically developing children, multisensory integration (MSI) abilities were known to continue improving late into childhood. The current study



looked at whether one aspect of MSI—integrating audio and visual speech signals—continues to develop in high-functioning children with ASD as well.

In the study, 222 children ages 5 to 17, including both typically developing children and high-functioning children with ASD, were tested for how well they could understand speech with increasing levels of background noise. In one test, the researchers played <u>audio recordings</u> of simple words. In a second test, the researchers played a video of the speaker articulating the words, but no audio. A third test presented the children with both the audio and video recordings.

The test mimics the so-called "cocktail party" effect: a noisy environment with many different people talking. In such settings, people naturally rely on both auditory and facial clues to understand what another person is saying. "You get a surprisingly big boost out of lipreading, compared with hearing alone," said Dr. Foxe. "It's an integrative process."

In the first test (audio alone), the children with ASD performed almost as well as typically developing children across all age groups and all background noise levels. In the second test (video alone), the children with ASD performed significantly worse than the typically developing children across all age groups and all background noise levels. "But the typically developing children didn't perform very well, either," said Dr. Foxe. "Most people are fairly terrible at lip-reading."

In the third test (audio and video), the younger children with ASD, ages 6 to 12, performed much worse than the typically developing children of the same age, particularly at higher levels of <u>background noise</u>. However, among the older children, there was no difference in performance between the typically developing and children with ASD.



"In adolescence, something amazing happens and the kids with ASD begin to perform like the typically developing kids," said Dr. Foxe. "At this point, we can't explain why. It may be a function of a physiological change in their brain or of interventions they've received, or both. That is something we need to explore."

The researchers acknowledge some limitations to their study. "Instead of doing a cross-sectional study like this, where we tested children at various ages, we would prefer to do a longitudinal study that would involve the same kids who'd be followed over the years from childhood through adolescence," Dr. Foxe said. "We also need to find a way to study what is happening with low- and mid-functioning children with ASD. They are much less tolerant of testing and thus harder to study."

According to the researchers, the work highlights the need to develop more effective therapies to help ASD <u>children</u> better integrate audio and visual speech signals. "We are beginning to work on that," said Dr. Foxe.

More information: The paper is titled "Severe Multisensory Speech Integration Deficits in High-Functioning School-Aged Children with an Autism Spectrum Disorder (ASD) and their Resolution during Early Adolescence."

Provided by Albert Einstein College of Medicine

Citation: Autistic children can outgrow difficulty understanding visual cues and sounds (2013, August 28) retrieved 5 May 2024 from <u>https://medicalxpress.com/news/2013-08-autistic-children-outgrow-difficulty-visual.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.