

High-flying pilots at increased risk of brain lesions

August 19 2013

A new study suggests that pilots who fly at high altitudes may be at an increased risk for brain lesions. The study is published in the August 20, 2013, print issue of *Neurology*, the medical journal of the American Academy of Neurology.

For the study, 102 U-2 United States Air Force pilots and 91 non-pilots between the ages of 26 and 50 underwent MRI brain scans. The scans measured the amount of [white matter hyperintensities](#), or tiny [brain lesions](#) associated with [memory decline](#) in other neurological diseases. The groups were matched for age, education and [health factors](#).

"Pilots who fly at altitudes above 18,000 feet are at risk for decompression sickness, a condition where gas or atmospheric pressure reaches lower levels than those within body tissues and forms bubbles," said study author Stephen McGuire, MD, with the University of Texas in San Antonio, the US Air Force School of Aerospace Medicine and a Fellow of the American Academy of Neurology. "The risk for decompression sickness among Air Force pilots has tripled from 2006, probably due to more frequent and longer periods of exposure for pilots. To date however, we have been unable to demonstrate any permanent clinical neurocognitive or memory decline."

Symptoms affecting the brain that sometimes accompany decompression sickness include slowed thought processes, confusion, unresponsiveness and permanent [memory loss](#).

The study found that pilots had nearly four times the volume and three times the number of brain lesions as non-pilots. The results were the same whether or not the pilots had a history of symptoms of decompression sickness.

The research also found that while the lesions in non-pilots were mainly found in the frontal white matter, as occurs in normal aging, lesions in the pilots were evenly distributed throughout the brain.

"These results may be valuable in assessing risk for occupations that include high-altitude mountain climbing, deep sea diving and high-altitude flying," McGuire said.

Provided by American Academy of Neurology

Citation: High-flying pilots at increased risk of brain lesions (2013, August 19) retrieved 23 April 2024 from <https://medicalxpress.com/news/2013-08-high-flying-brain-lesions.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|