

# Neurocognitive testing more accurate than self-reporting of cheerleaders' concussion recovery

August 8 2013

---

Concussions have become a major public health issue, with both short- and long-term side effects. In sports, cheerleading has the highest rate of catastrophic injury, with some studies reporting approximately 6% of total injuries as concussions. Return-to-play guidelines have relied on athletes' self-reports; however, this has led to concerns about the ability of athletes to truly recognize their own symptoms and recovery. In a new study scheduled for publication in *The Journal of Pediatrics*, researchers evaluate the accuracy of neurocognitive testing compared with self-reported symptoms of concussions in cheerleaders.

In the study, 138 junior and senior high school cheerleaders with concussions underwent pre-season baseline neurocognitive testing and completed at least one follow-up evaluation within 7 days of injury using Immediate Post-Concussion Assessment and Cognitive Testing (ImPACT). According to Dr. Mark R. Lovell and a colleague from Vanderbilt University School of Medicine, "We hypothesized that the use of ImPACT would result in an increased capacity to detect and measure post-concussive abnormalities in cheerleaders compared with symptom assessment alone." All concussions were diagnosed by a physician, athletic trainer, or other school health care official who was present at the time of injury.

Overall, 62% of the cheerleaders reported an increase in symptoms after a concussion (e.g., [headache](#), nausea, and [dizziness](#)) compared with their

baseline. Of the cheerleaders who denied an increase in concussion symptoms from baseline, 33% had at least one ImPACT score that exceeded index criteria. This means that these cheerleaders reported their symptoms inaccurately, overestimated their recovery, or were unaware of their decreased neurocognitive performance.

The results show that the addition of a neurocognitive assessment could be a useful tool to evaluate when cheerleaders with concussion have returned to normalized baseline measures. They also support the idea that self-reported symptoms and decreased neurocognitive test scores after concussion may differ. According to co-author Dr. Gary Solomon, "It is common knowledge that athletes may at times minimize or deny symptoms after injury to avoid being removed from competition." Treating physicians should be cautious to return athletes to play based solely on self-reported [symptoms](#).

**More information:** "Neurocognitive Test Performance and Symptom Reporting in Cheerleaders with Concussions," by Mark R. Lovell, Ph.D., and Gary S. Solomon, Ph.D., appears in *The Journal of Pediatrics*, [DOI: 10.1016/j.jpeds.2013.05.061](https://doi.org/10.1016/j.jpeds.2013.05.061)

Provided by Elsevier

Citation: Neurocognitive testing more accurate than self-reporting of cheerleaders' concussion recovery (2013, August 8) retrieved 19 April 2024 from <https://medicalxpress.com/news/2013-08-neurocognitive-accurate-self-reporting-cheerleaders-concussion.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--