

Prolactin reduces arthritis inflammation

August 1 2013

Inflammatory joint diseases such as rheumatoid arthritis are the result of cartilage damage and loss. Chondrocytes are the only cells that are found in cartilage and their death is linked to decreased cartilage health.

In this issue of the *Journal of Clinical Investigation*, Carmen Clapp and colleagues at the National University of Mexico identify prolactin as a potential treatment for inflammatory joint disease.

Prolactin treatment prevented chondrocyte death and associated cartilage degradation. In a rat model of inflammatory arthritis, prolactin treatment reduced inflammation, bone erosion, joint swelling, and pain.

This study indicates that prolactin therapy has the potential to relieve many symptoms associated with rheumatoid arthritis and other inflammatory-related diseases.

More information: Prolactin promotes cartilage survival and attenuates inflammation in inflammatory arthritis, *J Clin Invest*.
[doi:10.1172/JCI69485](https://doi.org/10.1172/JCI69485)

Provided by Journal of Clinical Investigation

Citation: Prolactin reduces arthritis inflammation (2013, August 1) retrieved 3 May 2024 from <https://medicalxpress.com/news/2013-08-prolactin-arthritis-inflammation.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.