

A vision exam for mice

August 21 2013

How can one use simple means to investigate the visual abilities of animals? This question is being pursued by the research group of Dr. Thomas Münch at the Centre for Integrative Neuroscience at the University of Tübingen. Dr. Münch is, among other things, interested in the restoration of lost visual function. Mice can suffer from similar diseases of blindness as humans, so that new therapeutic approaches can be developed by doing research on mice.

Since the treated animals cannot tell us how well they can see, one attempts to draw conclusions about visual abilities from behavioral observations. Animals are commonly trained with food to display a certain behavior in response to color or pattern. This experimental approach, however, is very time-consuming, and training blind animals is nearly impossible. The researchers therefore take advantage of a simple reflex: the optokinetic reflex. This reflex can easily be noticed on a train ride: when observing the landscape from a moving train, the eyes periodically follow the passing landscape. This effect keeps the image of [moving objects](#) constant on the [retina](#) for a short while. This phenomenon occurs in most animals.

A new study published this week in the journal *Behavioral Neuroscience* is based on this effect. The [visual performance](#) of mice was examined using a chamber whose walls were made of four computer screens. The monitors showed a rotating striped pattern. This simulated movement of the environment triggered the optokinetic reflex and the mouse followed the pattern with its gaze – but only when the animal could recognize the striped pattern. This allows the visual performance of each individual

animal to be determined. Just as the [ophthalmologist](#) can make the displayed pattern finer during vision testing, so too can the displayed pattern be changed until the animal is no longer able to recognize it and subsequently fails the reflex test.

The utilization of the [computer monitors](#) allows the contrast or resolution of the pattern to be changed in almost any way desired. For the study, doctoral candidate Boris Benkner has developed software that automatically evaluates the animal's behavior and can thus determine the visual ability of an animal in a short time. "In previous studies, it was necessary to tediously analyze each animal's behavior manually," said Benkner. "Our automated method is not only faster but also more objective, because the stripe pattern does also influence the observer when doing the analysis."

The senior author of the study, Thomas Münch, sees great potential in the newly developed method to evaluate new treatment strategies for blindness. "Currently, many new ways for treating blindness are being developed, ranging from dietary supplements to optogenetics to stem cell therapy" said Münch. "It is important to scrutinize these therapies from the beginning, to test if they really improve the [visual abilities](#) of the treated animals."

More information: Benkner, B. et al. Characterizing Visual Performance in Mice: An Objective and Automated System Based on the Optokinetic Reflex, *Behavioral Neuroscience*, Online First Publication, Aug. 20, 2013. [DOI: 10.1037/a0033944](https://doi.org/10.1037/a0033944)

Provided by University of Tübingen

Citation: A vision exam for mice (2013, August 21) retrieved 24 May 2024 from

<https://medicalxpress.com/news/2013-08-vision-exam-mice.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.