

# Biological therapy with cediranib improves survival in women with recurrent ovarian cancer

September 30 2013

---

Women with ovarian cancer that has recurred after chemotherapy have survived for longer after treatment with a biological therapy called cediranib, according to new results to be presented today (Monday) at the 2013 European Cancer Congress (ECC2013) [1].

Cediranib, which is taken in pill form, is an inhibitor of a cell signalling process involved in formation of tumour blood vessels, essential for tumour growth, and it is the first oral inhibitor of its kind to show an improvement in the time before patients' disease progresses and in overall survival. The drug is a tyrosine kinase inhibitor, a type of [biological therapy](#) that blocks vascular endothelial growth factor (VEGF) receptors, which control the development of blood vessels required for growing tumours.

Professor Jonathan Ledermann, Professor of Medical Oncology at UCL Cancer Institute, University College London, presented first results from ICON6, an international randomised, double-blind, academic clinical phase III trial of cediranib.

"In women whose [ovarian cancer](#) had been treated with platinum-based chemotherapy together with cediranib given during and after the chemotherapy, we found that the time before the tumour started to grow again was extended by an average of 3.2 months. This sounds like a modest increase but represents about a 30% improvement, with overall

survival also increased by a similar amount, to an average of 2.7 months over a two-year period of follow-up," he said.

Studies with chemotherapy alone have shown that the time before patients experience disease progression following treatment for a relapse that is sensitive to platinum-based chemotherapy is an average of eight to nine months. These latest results show that cediranib in addition to chemotherapy increased the time before the disease progressed from 9.4 to 12.6 months over a period of two years, and it extended overall survival from 17.6 to 20.3 months.

"These are ground-breaking data," said Prof Ledermann. "Cediranib is the first oral VEGF tyrosine kinase inhibitor that has been shown to delay tumour progression and improve overall survival in recurrent ovarian cancer. It is simple to give for a prolonged period and in most patients it is well-tolerated." Adverse side-effects included high blood pressure, diarrhoea and fatigue.

A total of 456 patients whose ovarian cancer had recurred were enrolled in the trial in 63 centres from the UK, Canada, Australasia and Spain. They were randomised to receive platinum-based chemotherapy together with a placebo (the reference arm of the trial), or 20 mg a day of cediranib during chemotherapy followed by placebo for 18 months (concurrent arm of the trial), or 20 mg a day of cediranib during chemotherapy followed by cediranib as a maintenance treatment (maintenance arm).

"ICON6 has a three-arm design in which the effect of cediranib given with chemotherapy and continued as maintenance can be compared with standard chemotherapy. This is the first trial to have demonstrated a benefit of concurrent cediranib with [chemotherapy](#), as well as demonstrating an additional benefit with maintenance cediranib," he said.

An increased survival time of nearly three months is significant in this group of patients. Prof Ledermann explained: "In previous ovarian cancer trials any improvement seen with each new treatment has been incremental. Survival has improved through sequential use of drugs. Most of the recent positive trials have shown an improvement in progression-free survival. Trials showing an improvement in overall survival are uncommon. Cediranib produces an incremental improvement in progression-free survival and an incremental improvement in overall survival. Although the average improvement in overall survival is 2.7 months, some patients will see a much more substantial benefit."

ECCO president, Professor Cornelis van de Velde, commented: "These are important results for women with recurrent ovarian cancer. Once the disease has recurred there are few treatment options available that make a significant difference to its progression and to overall survival. The ICON6 trial shows that cediranib does make a difference and it is to be hoped that it can be made available to women as soon as is practicable."

**More information:** [1] The 2013 European Cancer Congress is the 17th congress of the European Cancer Organisation (ECCO), the 38th congress of the European Society for Medical Oncology (ESMO) and the 32nd congress of European Society for Therapeutic Radiology and Oncology (ESTRO).

[2] ICON6 is an academic trial run by the UK Medical Research Council's Clinical Trials Unit, funded by Cancer Research UK, and supported by AstraZeneca. It is a Gynecologic Cancer Intergroup (GCIG) trial with participating groups from UK (NCRI and SGCTG), Canada (NCIC Clinical Trials Group), Australia/New Zealand (ANZGOG) and Spain (GEICO).

[3] Cediranib is an investigational compound manufactured by

AstraZeneca and is not currently approved for use in the European Union or United States.

Provided by The European Cancer Organisation (ECCO)

Citation: Biological therapy with cediranib improves survival in women with recurrent ovarian cancer (2013, September 30) retrieved 27 April 2024 from <https://medicalxpress.com/news/2013-09-biological-therapy-cediranib-survival-women.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.