

Study shows colonoscopy better than sigmoidoscopy in protecting against colorectal cancer

September 18 2013

A study in the Sept. 19 *New England Journal of Medicine* finds that colonoscopy appears to reduce the risk of developing or dying from colorectal cancer more powerfully than does sigmoidoscopy, a similar procedure that examines only a portion of the colon. The investigation, which analyzes data from two long-term studies, also identifies molecular features that may help explain tumors that are diagnosed despite an individual's having recently undergone colonoscopy.

"This study provides some of the clearest evidence to date that colonoscopy has advantages over sigmoidoscopy for the prevention of colorectal [cancer](#), although it is not a perfect test," says Andrew Chan, MD, MPH, of the Massachusetts General Hospital (MGH) Gastrointestinal Unit, corresponding and co-senior author of the report. "Our findings support current recommendations for colorectal cancer screening with either of these procedures and provide a rationale for some individuals to consider colonoscopy as a preferred test for their individual situation."

Both colonoscopy and sigmoidoscopy allow examination of the internal surface of the colon through a fiberoptic tube with a light and camera at the end, but while colonoscopy visualizes the entire colon, sigmoidoscopy only reaches the lower third on the left side of the body. Current recommendations for colorectal screening in average-risk individuals over age 50 are either a sigmoidoscopy every 5 years –

followed by colonoscopy if abnormalities are detected – or a colonoscopy every 10 years.

While randomized, controlled [clinical trials](#) have confirmed that sigmoidoscopy reduces the incidence of and death from colorectal cancer, similar trials have not yet been done for colonoscopy, and some prior studies have suggested that colonoscopy did not have significant advantages over sigmoidoscopy. However, the more comprehensive colonoscopy is widely recommended for colorectal screening based on its ability to visualize the whole colon. The current study was designed to compare the effectiveness of both procedures in reducing the long-term risk of colorectal cancers in different segments of the colon and to examine molecular features of tumors developing soon after a negative examination.

To do this, the investigators analyzed information from almost 89,000 participants in the Nurses Health Study and the Health Professionals Follow-up Study, both of which gather comprehensive health information from their participants every two years. From 1988 to 2008, the participants reported whether they had either sigmoidoscopy or colonoscopy in the two preceding years and, if they had, whether the examinations were for screening purposes or because of symptoms like bleeding. For participants who developed cancer or who reported having colon polyps removed, the researchers obtained consent to review their medical records and pathology reports.

Over the 22 years for which data was analyzed, 1,815 cases of colorectal cancer and 474 related deaths were documented in the 88,902 study participants. While a negative result – meaning that neither cancer nor polyps were diagnosed – in either examination or having polyps that were removed was associated with a reduced risk of developing left-side colon cancer, only a negative colonoscopy was associated with significant risk reduction in the entire colon.

Compared with the 15 percent of study participants who had no endoscopic colon exam during the study period, having either procedure significantly reduced the overall risk of dying from colorectal cancer. But while sigmoidoscopy only cut the risk of dying from left-side tumors, screening colonoscopy reduced the risk of any colorectal-cancer-associated death.

Having a single negative colonoscopy significantly reduced the incidence of any colorectal cancer for up to 15 subsequent years, supporting the current recommendations for a 10-year interval between exams. Among participants who had polyps removed, the incidence of subsequent tumors was reduced for up to 5 years, except for those whose polyps had features indicating elevated risk, in whom the risk reduction was not as strong.

Among the participants who did develop colorectal cancer, molecular data was available on 62 tumors that had been diagnosed within 5 years after colonoscopy. The researchers identified specific cancer-related features more likely to be seen in those tumors than in cancers diagnosed after more time had passed. "It is widely believed that tumors with those features develop from a specific type of polyp called a sessile serrated adenoma, which may be particularly difficult to detect endoscopically or to remove completely," Chan explains. "Focused efforts to improve detection and removal of these polyps are a high priority.

"For individual patients, the decision to select colonoscopy over [sigmoidoscopy](#) for colorectal [cancer screening](#) requires discussing with their physicians how the advantages and disadvantages of each test apply to their personal situations," he adds. "Right now our data support the use of colonoscopy if the primary goal is maximum risk reduction for both left- and right-side colorectal cancers. Future study should focus on ways of further improving [colonoscopy](#) quality, as well as better understanding why some tumors arise within a short time after a negative

examination." Chan is an associate professor of Medicine at Harvard Medical School.

Provided by Massachusetts General Hospital

Citation: Study shows colonoscopy better than sigmoidoscopy in protecting against colorectal cancer (2013, September 18) retrieved 19 April 2024 from

<https://medicalxpress.com/news/2013-09-colonoscopy-sigmoidoscopy-colorectal-cancer.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--