

Prenatal exposure to antiepileptic drugs associated with impaired fine motor skills

September 23 2013

Prenatal exposure to antiepileptic medications was associated with an increased risk of impaired fine motor skills (small muscle movements) in children at age 6 months, but breastfeeding by women taking the medications was not associated with any harmful effects on child development at ages 6 to 36 months, according to a report published by *JAMA Neurology*.

Few studies have examined development during the first months of life of [children](#) of mothers with epilepsy, according to background in the study by Gyri Veiby, M.D., of the University of Bergen, Norway, and colleagues.

Researchers used data from the Norwegian Mother and Child Cohort Study from 1999 to 2009, in which mothers reported children's motor and social skills, language and behavior at 6, 18 and 36 months of age. Women also provided information on breastfeeding during the first year.

Exposure to antiepileptic drugs during pregnancy was reported in 223 children, mostly to a single drug (n=182).

At age 6 months, a higher proportion of infants whose mothers took antiepileptic drugs had impaired fine [motor skills](#) compared to the reference group (11.5 percent vs. 4.8 percent). The use of multiple antiepileptic medications was associated with impaired fine motor and social skills, according to the study results.

Breastfeeding by mothers using antiepileptic drugs was not associated with adverse development at ages 6 to 36 months in nursing children, according to the study.

"Women with epilepsy should be encouraged to breastfeed their children irrespective of antiepileptic medication use," the study concludes.

In an editorial, Paul C. Van Ness, M.D., of the University of Texas Southwestern Medical Center, Dallas, writes: "Pregnant women with epilepsy often ask whether they will be able to breastfeed. Many have been given conflicting advice when there were scant data to answer the question."

"This study adds additional evidence that long-term breastfeeding is safe and perhaps even beneficial to infants of [mothers](#) taking AEDs [[antiepileptic drugs](#)]," Van Ness continues.

"This should be an important area of discussion with any woman of childbearing potential well in advance of pregnancy using the best available information. Thus, a woman should take folic acid supplements, received AED monotherapy whenever possible, and take the lowest effective dose of medication," the editorial concludes.

More information: *JAMA Neurol.* Published online September 23, 2013. [DOI: 10.1001/jamaneurol.2013.4290](https://doi.org/10.1001/jamaneurol.2013.4290)
JAMA Neurol. Published online September 23, 2013. [DOI: 10.1001/jamaneurol.2013.4348](https://doi.org/10.1001/jamaneurol.2013.4348)

Provided by The JAMA Network Journals

Citation: Prenatal exposure to antiepileptic drugs associated with impaired fine motor skills

(2013, September 23) retrieved 1 May 2024 from

<https://medicalxpress.com/news/2013-09-prenatal-exposure-antiepileptic-drugs-impaired.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.